

The European Association for Endoscopic Surgery clinical practice guideline on the pneumoperitoneum for laparoscopic surgery

J. Neudecker,^{1,2,3} S. Sauerland,^{1,3} E. Neugebauer,^{1,3} R. Bergamaschi,⁴ H. J. Bonjer,⁵ A. Cuschieri,⁶ K.-H. Fuchs,⁷ Ch. Jacobi,¹ F. W. Jansen,⁸ A.-M. Koivusalo,⁹ A. Lacy,¹⁰ M. J. McMahon,¹¹ B. Millat,¹² W. Schwenk¹

¹ Conference Organizers on behalf of the Scientific Committee of the European Association for Endoscopic Surgery (E.A.E.S.), c/o E.A.E.S. Office, Post Office Box 335, 5500 AH Veldhoven, The Netherlands

² Department of Surgery, Charité Campus Mitte, Humboldt-University of Berlin, Schumannstraße 20/21, 10117 Berlin, Germany

³ Biochemical and Experimental Division, 2nd Department of Surgery, University of Cologne, Ostmerheimer Strasse 200, 51109 Cologne, Germany

⁴ Department of Surgery, SSSF Hospital, University of Bergen, 6800 Førde, Norway

⁵ Department of Surgery, University Hospital Dijkzigt, Dr. Molewaterplein 40, 3015 GD Rotterdam, The Netherlands

⁶ University Department of Surgery, Ninewells Hospital and Medical School, Dundee DD1 9SY, United Kingdom

⁷ Department of Surgery, University of Würzburg, Josef-Schneider-Str. 2, 97080 Würzburg, Germany

⁸ Department of Gynecology, Leiden University, Post Office Box 9600, 2300 RC Leiden, The Netherlands

⁹ Department of Anaesthesia, University of Helsinki, 00130 Helsinki, Finland

¹⁰ Department of Surgery, Hospital Clinic, Villarroel 170, 08036 Barcelona, Spain

¹¹ Leeds Institute for Minimally Invasive Therapy (LIMIT), Great George Street, Leeds LS1 3EX, United Kingdom

¹² Department of Surgery, Hospital Saint Eloi, Avenue Bertin Sans No. 2, 34295 Montpellier, France

Received: 17 September 2001/Accepted in final form: 26 September 2001/Online publication: 20 May 2002

Abstract

Background: The pneumoperitoneum is the crucial element in laparoscopic surgery. Different clinical problems are associated with this procedure, which has led to various modifications of the technique. The aim of this guideline is to define the scientifically proven standards of the pneumoperitoneum.

Methods: Based on systematic literature searches (Medline, Embase, and Cochrane), an expert panel consensually formulated clinical recommendations, which were graded according to the strength of available literature evidence.

Recommendations: Preoperatively, all patients should be assessed for the presence of cardiac, pulmonary, hepatic, renal, or vascular comorbidity. Presupposing appropriate perioperative measures and surgical technique, there is no reason to contraindicate pneumoperitoneum in patients with peritonitis or intraabdominal malignancy. During laparoscopy, monitoring of end tidal CO₂ concentration is mandatory. The available data on closed- (Veress needle) and open-access techniques do not allow us to principally favor the use of either technique. Using 2 to 5-mm instead of 5 to 10-mm trocars improves

cosmetic result and postoperative pain marginally. It is recommended to use the lowest intraabdominal pressure allowing adequate exposure of the operative field, rather than using a routine pressure. In patients with limited cardiac, pulmonary, or renal function, abdominal wall lifting combined with low-pressure pneumoperitoneum might be an alternative. Abdominal wall lifting devices have no clinically relevant advantages compared to low-pressure (5–7 mmHg) pneumoperitoneum. In patients with cardiopulmonary diseases, intra- and postoperative arterial blood gas monitoring is recommended. The clinical benefits of warmed, humidified insufflation gas are minor and contradictory. Intraoperative sequential intermittent pneumatic compression of the lower extremities is recommended for all prolonged laparoscopic procedures. For the prevention of postoperative pain a wide range of treatment options exists. Although all these options seem to reduce pain, the data currently do not justify a general recommendation.

Only 15 years after the introduction of laparoscopic cholecystectomy, laparoscopic techniques (used either as a diagnostic tool or as a therapeutic access method) are among the most common procedures in surgery worldwide. However, concerns about higher surgical complications rates (such as vascular and intestinal injuries) compared to conventional techniques and anesthetics

Table 1. A method for grading recommendations according to scientific evidence^a

Grade of recommendation	Level of evidence	Possible study designs for the evaluation of therapeutic interventions
A	1a	Systematic review (with homogeneity) of RCTs
	1b	Individual RCT (with narrow confidence interval)
	1c	All-or-none case series
B	2a	Systematic review (with homogeneity) of cohort studies
	2b	Individual cohort study (including low-quality RCT)
	2c	“Outcomes” research
	3a	Systematic review (with homogeneity) of case-control studies
	3b	Individual case-control study
C	4	Case series (and poor-quality cohort and case-control studies)
D	5	Expert opinion without explicitly critical appraisal, or based on physiology, bench research, or “first principles”

^a From Sackett et al. [255]

logical risks have remained. Since the start of the laparoscopic era, numerous studies have described pathophysiological or clinical problems that are related to laparoscopy. Therefore, many technical innovations and modifications have been developed to improve safety and effectiveness of laparoscopy, but not all of them have been studied adequately before clinical use.

With these developments in mind, the European Association for Endoscopic Surgery (E.A.E.S.) decided to develop authoritative and evidence-based clinical practice guidelines on the pneumoperitoneum and its sequelae. The scope of these guidelines covers all important general surgical aspects of the pneumoperitoneum but not special laparoscopic procedures for defined pathologies. They address the pathophysiological basis for the clinical indications, aspects to establish the pneumoperitoneum, and perioperative aspects such as adhesions and pain. In addition, a clinical algorithm was formulated for practical use.

Methods

Under the mandate of the E.A.E.S Scientific Committee with the aim to set up evidence-based clinical practice guidelines, we combined the methodologies of a systematic review and a consensus development conference (CDC) because previous CDCs (both within and outside the E.A.E.S) had difficulties in identifying all relevant articles [218, 262, 280]. As a framework of the process, the key aspects pertaining to the pneumoperitoneum were precisely formulated in separate questions, which then were answered concurrently by the use of literature and expert evidence.

For the systematic review, one researcher (J.N.) performed comprehensive literature searches in Medline, Embase, and the Cochrane Library. We used the medical subject headings Laparoscopy and Pneumoperitoneum. Our primary intention was to identify all clinically relevant randomized controlled trials (RCTs). However, other trials using concurrent cohorts (CCTs), external or historical cohorts, population-based outcomes studies, case series, and case reports were accepted for a comprehensive evaluation of the pneumoperitoneum and its sequelae (Table 1). Included articles were scrutinized and classified by two reviewers (J.N. and S.S.). Furthermore, all panelists were asked to search the literature according to a list of defined questions. The reference lists of all relevant articles were also checked.

For the CDC, the conference organizers in Cologne, together with the scientific committee of the EAES, nominated a multidisciplinary expert panel. The criteria for selection were clinical and scientific expertise in the field of laparoscopy and geographical location within Europe.

Six months before the conference, the questions on laparoscopy were sent to the panelists. In parallel, the questions were answered by literature evidence found in systematic searches. One month before the conference, all answers from the panel and the literature searches were analyzed and subsequently combined into a provisional preconsensus statement and a clinical algorithm. Each panel member was also informed about the identities of the other members, which had not been previously disclosed.

In Maastricht, all panelists (except A.C. and H.J.B.) met for a first meeting on June 13, 2001. Here, the provisional bottom-line statements typed in bold and the clinical algorithm with the grades of recommendation were scrutinized word by word in a 5-h session in a nominal group process. For all statements, internal (expert opinion) and external evidence was compared. The following day the modified statement and the algorithm were presented to the conference audience by all panelists for public discussion (1.5-h session). During a postconsensus meeting on the same day, all suggestions from the audience were discussed again by the panelists, and the statement was further modified. The finalized statement as given later was mailed to all panelists for final approval (Delphi process) before duplication.

To increase readability, a short version of the clinical practice guidelines with a clinical algorithm was prepared (Fig. 1). The extended version consists of a detailed appraisal of pathophysiologic background and clinical research evidence. Each recommendation is graded according to its reliability and the rigor of research evidence behind the statement (Table 1).

Pathophysiological basis for the clinical indications

Cardiovascular system

Cardiovascular effects of pneumoperitoneum occur most often during its induction, and this should be considered when initial pressure is increased for introduction of access devices. In ASA I and II patients, the hemodynamic and circulatory effects of a 12- to 14-mmHg capnoperitoneum are generally not clinically relevant (grade A). Due to the hemodynamic changes in ASA III and IV patients, however, invasive measurement of blood pressure or circulating volume should be considered (grade A). These patients should also receive adequate preoperative volume loading (grade A), beta-blockers (grade A), and intermittent sequential pneumatic compression of the lower limbs, especially in prolonged laparoscopic procedures (grade C). If technically feasible, gasless or low-pressure laparoscopy might be an alternative for patients with limited cardiac function (grade B). The use of other gases (e.g., helium) showed no clinically relevant hemodynamic advantages (grade A).

Preoperative assessment

Patient is scheduled for laparoscopic surgery

Define patient as comorbid if:
ASA III-IV, COPD, NYHA III-IV,
CRF etc. (see text for details)

Pre-surgery interventions

Administer adequate preoperative volume loading (grade A)

Is patient comorbid ?

Yes

- Start invasive measurement of blood pressure or circulating volume (grade A)
- Insert urine catheter (grade B)
- Consider pharmacological interventions (e.g. β -blocker, nitroglycerin) (grade A)
- Consider gasless laparoscopy (grade B)
- In COPD patients, consider helium as alternative gas (grade B)

No

Start monitoring of end tidal CO₂-concentrations (grade A)

Is patient comorbid ?

Yes

- Use intermittent pneumatic compression (grade C)
- Use external heating devices

No

Estimated duration of surgery >2h ?

Yes

No

Surgical interventions

Establish pneumoperitoneum either by open or closed access technique (grade A)

Apply lowest possible pressure level (grade A)

Use small instruments, if suitable (grade A)

Perform surgery

At the end of operation, remove residual gas (grade B)

Fig. 1. Evidence-based clinical algorithm on the pneumoperitoneum for laparoscopic surgery. The recommendation is graded according to Table 1. Diamond boxes, decision boxes, square boxes action boxes [255].

Pneumoperitoneum decreases venous return, preload, and cardiac output (CO) and increases heart rate (HR), mean arterial pressure (MAP), as well as systemic (SVR) and pulmonary vascular resistance (PVR). These hemodynamic and cardiovascular - changes mostly occur because of increased intraabdominal pressure (IAP) (1b [159, 221, 291]) and the stimulated neurohumoral vasoactive systems [vasopressin and renin-aldosterone-angiotensin system (RAAS)] (1b [142, 158]), but are independent of type of gas (1b [28]). However, in oth-

erwise healthy patients these changes are not dangerous when IAP does not exceed 15 mmHg (1b [27]).

Increased IAP, up to 12–15 mmHg, decreases venous return, which results in reduced preload and CO, without adequate intravascular volume loading (1b [63, 142, 162, 201, 221]). Additionally, changes in body position, especially head-up tilt position, intensify these negative effects of a pneumoperitoneum (2b [115, 116]), whereas head-down or Trendelenburg position has a positive effect on venous return (1b [162]). Furthermore, the use

of positive end-expiratory pressure (PEEP) of 10 H₂O during pneumoperitoneum decreases preload and CO (4 [164]). Pneumoperitoneum increases sympathetic cardiac activity (1b [260]) and induces a hemodynamic stress response by activation of the neurohumoral vasoactive system (i.e., vasopressin and RAAS) resulting in increased HR, increased SVR and PVR, and increased arterial blood pressure (1b [142, 159]). This stress response leads to an increase in oxygen consumption, which might be deleterious for patients with compromised cardiac function. In clinical studies on ASA III and IV patients distinct intraoperative hemodynamic changes during pneumoperitoneum were described (4 [127]), but cardiovascular stability was unimpaired (4 [64, 83, 111, 322]) if appropriate invasive monitoring and pharmacologic interventions were used (4 [79, 292]). In contrast, there are reports of cardiovascular alterations persisting after release of the pneumoperitoneum (4 [108]). Most of these studies used an IAP of 12–15 mmHg without preoperatively volume loading. Without adequate intravascular volume loading a pneumoperitoneum in connection with head-up tilt position decreases CO significantly (up to 50%) (1b [142, 221]). In comorbid patients (ASA III and IV), RCTs with adequate sample size are missing.

In the majority of patients (ASA I and II), the hemodynamic effects of a pneumoperitoneum are without consequences and vanish after desufflation. Therefore, most patients without comorbidities (ASA I and II) do not need invasive hemodynamic monitoring. However, in ASA III and IV patients an invasive monitoring of blood pressure and circulating volume must be considered because only these measures allow early recognition and adequate treatment of severe cardiovascular changes (1b [162]). For intraoperative monitoring, a pulmonary artery catheter or COLD (cardiac oxygenation and lung water determination) monitoring should be applied, because transesophageal echocardiography in patients with cardiac disease has not been proven to be useful (4 [241]). For patients with severely compromised circulation, measurement of the pulmonary artery pressure (PAP) and CO can be necessary. However, interpreting changes in central venous pressure (CVP) and PAP may be difficult (4 [213]). Due to the consecutive increase in intrathoracic pressure during laparoscopy CVP and PAP also increase, but right arterial volume is not decreased. Therefore, CVP may only incorrectly describe the effective circulating blood volume and could be misinterpreted [162].

Since the effects of increased IAP on hemodynamics are volume dependent, adequate preoperative intravascular loading is essential, especially in patients with cardiac diseases, to prevent cardiovascular side effects of a pneumoperitoneum (1b [162]). Another intervention of proven effectiveness that also increases cardiac preload and thereby prevents hemodynamic changes (2b [6]) is intermittent sequential pneumatic compression of the lower extremities to augment venous blood return (1b [273, 274], 2b [273, 274]).

To minimize the effects of hemodynamic stress response on myocardial oxygen consumption, esmolol or clonidine can safely be used (1b [142, 163]) if volume

depletion is not present. Intraoperative hemodynamic alterations in patients with underlying cardiopulmonary disease can be effectively controlled by appropriate pharmacological intervention (use of intravenous nitroglycerin) (2b [80]).

Hemodynamic and circulatory changes are independent from the used gas (CO₂ or helium) (1b [28]) but decreased during gasless laparoscopy (1b [5, 91, 159, 201, 221]). Therefore, gasless laparoscopy might be an alternative for patients with limited cardiac function. In summary, cardiac diseases are associated with an increased risk of general complications after laparoscopic surgery (and even higher risk after conventional surgery). Since various surgical and nonsurgical treatment options can be recommended to reduce these risks, the presence of heart disease does not principally contraindicate laparoscopic surgery (2b [239, 240]). There is a need for further trials in ASA III and IV patients.

Lung physiology and gas exchange

Carbon dioxide pneumoperitoneum causes hypercapnia and respiratory acidosis. During laparoscopy, monitoring of end-tidal CO₂ concentration is mandatory (grade A) and minute volume of ventilation should be increased in order to maintain normocapnia. Increased intraabdominal pressure and head-down position reduce pulmonary compliance and lead to ventilation--perfusion mismatch (grade A). In patients with normal lung function, these intraoperative respiratory changes are usually not clinically relevant (grade A). In patients with limited pulmonary reserves, capnoperitoneum carries an increased risk of CO₂ retention, especially in the postoperative period (grade A). In patients with cardiopulmonary diseases, intra- and postoperative arterial blood gas monitoring is recommended (grade A). Lowering intraabdominal pressure and controlling hyperventilation reduce respiratory acidosis during pneumoperitoneum (grade A). Gasless laparoscopy, low-pressure capnoperitoneum, or the use of helium might be alternatives for patients with limited pulmonary function (grade B). Laparoscopic surgery preserves postoperative pulmonary function better than open surgery (grade A).

The specifics of a capnoperitoneum, the IAP and the used gas, result in hypercapnia, respiratory acidosis, reduced pulmonary compliance, and increased airway resistance (1b [224, 307]). Additional changes in body position have minor influences (2b [68]) but could intensify these effects, especially in the head-down position (2b [114, 116]). Relaxation of the diaphragm caused by anesthesia in combination with increased intraabdominal pressure impairs excursion of the diaphragm and leads to compression of the lower lung lobes. These effects result in a decreased tidal volume, ventilation--perfusion mismatch, a decreased shunt volume, increased dead space, and decreased pulmonary compliance (1b [160], 2b [250]). Pulmonary gas exchange during laparoscopy can be optimized by the choice of the anesthetic procedure (1b [93]) and PEEP (5 [183]). Without hyperventilation $P_a\text{CO}_2$ will increase by 8–10 mmHg and pH will decrease (1b [315]) before a steady

state is reached. Intraoperatively, pulmonary changes due to capnoperitoneum are compensated by otherwise healthy adults (1b [197]).

Various laparoscopic procedures have been shown to result in better postoperative pulmonary function when compared to their open-surgery counterparts (1b [38, 42, 48, 51, 74, 112, 149, 167, 197, 206, 275, 307]), but clinically more relevant end points such as postoperative pulmonary complications were rarely evaluated or found unchanged in ASA I and II patients (1b [205, 206]). These data generally prove that laparoscopy rather than conventional surgery should be advised for compromised patients, particularly those with obstructive lung disease. This superiority of laparoscopic surgery during the postoperative period is mostly related to its lesser extent of surgical trauma and pain, but laparoscopy has certain effects on ventilation that deserve special attention.

Capnoperitoneum with an IAP of more than 12 mmHg combined with head-down or Trendelenburg position should be avoided because it reduces pulmonary compliance by more than 30% and there is ventilation-perfusion impairment (1b [155, 159, 224] 2b [188, 250, 296]). Hypercapnia and respiratory acidosis can be avoided by controlled hyperventilation (1b [315]). CO₂ storage during pneumoperitoneum can result in postoperative hypercapnic hangover, which has to be particularly considered in cases, of accidental subcutaneous CO₂ insufflation. To recognize these changes intra- and postoperative arterial blood gas monitoring and continuous capnometry are generally recommended for comorbid patients [161], particularly those with cardiopulmonary diseases (2b [314]). These patients may also benefit from prolonged postoperative mechanical ventilation [160]. From a more surgical standpoint, gasless laparoscopy or the use of other gases (e.g., helium and N₂O) may have clinically relevant advantages (1b [28, 155, 159, 224]), but the results of randomized trials are inconsistent (1b [92]) and need to be confirmed. Trocar positioning also has a relevant influence on pulmonary function, (1a [167]). Overall, most of the discussed randomized trials included only small numbers of patients, leading to an increased chance of type II error.

It should be mentioned that capnothorax can be a serious, albeit rare, complication that has been encountered in patients after capnoperitoneum (4 [7, 233]). Capnothorax occurs more often after laparoscopic esophageal or gastric surgery but has also been observed after lower abdominal procedures or even hernia repair. Because of the high solubility of CO₂, asymptomatic capnothorax diagnosed by postoperative chest x-ray may be treated conservatively. However, tension capnothorax may occur very rarely. Therefore, symptomatic capnothorax requires immediate drainage.

Venous blood return

During laparoscopy, both head-up position and elevated intraabdominal pressure independently reduce venous

blood return from the lower extremities (grade A). Intraoperative sequential intermittent pneumatic compression of the lower extremities effectively reduces venous stasis during pneumoperitoneum (grade A/B) and is recommended for all prolonged laparoscopic procedures. The incidence of thromboembolic complications after pneumoperitoneum is not known.

Increased intraabdominal pressure together with reverse Trendelenburg position (head-up position) decreases venous return from the lower extremities by more than 40% (1b [273, 274], 2b [12, 99, 123, 204]) with a concomitant increase in femoral venous pressure (2b [12, 139]). However, it has been hypothesized that the systemic coagulation system is activated by laparoscopic surgery (1b [243], 2b [41, 178]), but controversial data exist (2b [67, 119, 192]). Due to the impairment of lower extremity circulation, the increased venous pressure, and the activation of the systemic fibrinolytic system, the potential risk for deep venous thrombosis (DVT) is increased. Although there are alarming reports about a high incidence of DVT after pneumoperitoneum (4 [230]), the rate of clinically evident postoperative thromboembolic complications after laparoscopic surgery remains unclear [10, 31, 184, 189]. The negative effects of elevated IAP and body position on venous blood return from the lower extremities can partly be counteracted by intermittent sequential compression of the lower limbs in laparoscopic cholecystectomy and colorectal surgery (1b [273, 274], 2b [273, 274]). Whether such compression does reduce thromboembolic event remains to be elucidated in larger trials.

The effects of a low-pressure pneumoperitoneum (5–7 mmHg) and abdominal wall lifting on thromboembolic complications have not been studied, although from a pathophysiologic standpoint a positive effect can be reasonably anticipated.

Perfusion of intraabdominal organs

Although in healthy subjects (ASA I and II), changes in kidney or liver perfusion (grade A) and also splanchnic perfusion (grade D) due to an intraabdominal pressure of 12–14 mmHg have no clinically relevant effects on organ function, this may not be the case in patients with already impaired perfusion. Especially in patients with impaired hepatic or renal function or atherosclerosis, intraabdominal pressure should be as low as possible to reduce microcirculatory disturbances (grade B). Patients with impaired renal function should be adequately volume loaded before and during elevated intraabdominal pressure (grade A).

Renal effects

Randomized clinical trials showed a decrease in renal blood flow (RBF), glomerular filtration rate, and urine output in the initial phase of a pneumoperitoneum (1b [155, 221]). With increasing IAP renal function is gradually depressed (5 [146]). Elevated IAP causes renal

dysfunction due to direct mechanical compression of renal parenchyma, renal arteries, and veins (5 [247]). The reduction in RBF and urine output is probably caused by a decrease in CO and/or the compression of the renal vein. In experimental studies, renal vein flow remained decreased for at least 2 h postoperatively (5 [195, 247]). Mediated by humoral factors, a sympathetic reaction induces a constriction of the renal artery. Pneumoperitoneum increases plasma renin activity (PRA) and consequently activates the RAAS, which promotes the renal vasoconstriction via angiotensin II. However, one prospective randomized trial found no signs of a clinically relevant impairment of renal function (1b [26]).

Hepatoportal effects

When measured with laser Doppler, hepatoportal circulation is gradually decreased with increasing IAP (2b [69], 4 [136, 223]). In elderly patients, splanchnic circulation is very sensitive to elevated pressure (4 [261]). Experimental and clinical studies reported elevated liver enzymes after prolonged laparoscopic procedures and elevated intraabdominal pressure (1b [95], 2b [209]). However, in one RCT no signs of clinically relevant postoperative liver dysfunction were detected (1b [26]).

Splanchnic effects

Elevated IAP mechanically compresses capillary beds, decreases splanchnic microcirculation, and thus impairs oxygen delivery to the intraabdominal organs. During pneumoperitoneum, a 24% reduction of blood flow in the superior mesenteric artery and the hepatic portal vein was reported (5 [125]). In healthy patients, a high vs low IAP (15 vs 10 mmHg) decreased blood flow into the stomach (54%), the jejunum (32%), the colon (4%), the parietal peritoneum (60%), and the duodenum (11%) (4 [266]). Furthermore, clinical and animal studies noted a decrease in gastric intramucosal pH (1b [157], 2b [69]), which may be the earliest indicator of altered hemodynamic function compared to traditional measurements, such as CO, SVR, and lactate [154], but conflicting findings exist ([69, 187, 223]). The clinical implications of these investigations remain unclear.

Otherwise healthy patients seem to compensate changes in intraabdominal organ perfusion without impairment of organ function. However, in patients with cardiovascular comorbidities or preexisting organ disorders, reduced alteration in organ perfusion could have detrimental effects. Therefore, for these patients careful observation and selection of surgical technique are required.

Several studies of different quality reported that in patients with limited hepatic or renal function, postoperative hepatic and renal function were better preserved by keeping IAP under 12 mmHg and by avoiding a prolonged pneumoperitoneum (1b–4 [69, 125, 154, 266]). Recently, one experimental study investigated the influence of different IAP levels on intra- and extraabdominal tissue blood flow by using color-labeled micr-

spheres and reported, a nonimpaired tissue blood flow during capnoperitoneum of 10–12 mmHg (5 [317]). Esmolol inhibits the release of renin and blunts the pressor response to induction and maintenance of pneumoperitoneum. It may protect against renal ischemia during laparoscopy because urine output under, esmolol therapy was found to be higher (1b [162]). Nonsteroidal antiinflammatory drugs (NSAIDs), widely used in laparoscopic surgery, can cause renal medullary vasoconstriction. Because cases of renal failure after laparoscopic surgery and NSAID therapy were reported, NSAIDs should be replaced by other analgetics wherever possible (5; A.-M. Koivusalo, personal communication).

Stress response and immunologic parameters

Changes in systemic inflammatory and antiinflammatory parameters (mainly cytokines) as well as in stress response parameters are less pronounced after laparoscopic surgery than after conventional surgery (grade A). Whether this leads to clinically relevant effects (e.g., less pain, fatigue, and complications) remains to be proven. There is no compelling clinical evidence that specific modifications of the pneumoperitoneum alter the immunological response.

The influence of pneumoperitoneum on the function of the immune system and stress response is poorly evaluated because most studies investigate surrogate parameters of the immunological function, such as cytokines and other cell products, and not the cell function itself (e.g., count, ratio, concentration, and activity of immunological cells). The essential clinical outcomes after surgery concerning immunological functions are infections (e.g., sepsis, pneumonia, urinary tract infection, and local wound-related infections) and cancer growth (e.g., metastasis and local tumor spread). However, there is no study in the field of laparoscopic surgery demonstrating an association between changes of intra- and postoperative immune function and the occurrence of clinical complications.

Clinical controlled trials of laparoscopic versus conventional surgery have mostly focused on changes of cytokine levels to describe the influence of pneumoperitoneum on systemic immunological functions. These studies showed differences in serum cytokine levels between laparoscopic and conventional surgery for IL-1 (1b [174]), IL-6 (1b [36, 135, 165, 176, 254, 320, 322]), CRP (1b [135, 140, 176, 235, 254, 320]), CRP (1b [133, 138, 174, 233, 252, 318]) and cell-mediated immunity (1b [224]) that have not been confirmed by other authors (1b [17, 198], 2b [89]). In RCTs, postoperative immunological functions seemed to be better preserved after laparoscopic compared to conventional procedures (1b [13, 45, 151, 176, 235, 276, 284, 308, 321]); however, some trials found no differences (1b [73, 113, 173, 203, 226, 248, 270, 289, 295]) and one trial even reported a more pronounced immunodepression after laparoscopy (1b [290]). Additional RCTs examined perioperative stress response and found adrenaline (1b [150]), noradrenaline (1b [150]), and cortisol (1b [150, 174, 303]) decreased to a lesser extent after laparoscopic than after conventional

Table 2. Randomized clinical trials of Veress needle or open approach

Reference/year	No. of patients	Procedure	Access time (min)	Complications	Results
Gullà et al. [103]/2000	262	Diagnostic and operative laparoscopy	Not mentioned	Needle: 11/101 Open: 0/161	Open technique is safer
Saunders et al. [262]/1998	176	Diagnostic laparoscopy in abdominal trauma	Needle: 2.7 Open: 7.3	Needle: 0/98 Open: 0/78	Veress technique is faster
Cogliandolo et al. [50]/1998	150	Laparoscopic cholecystectomy	Needle: 4.5 Open: 3.2	Needle: 5/75 Open: 5/75	Open technique is faster
Peitgen et al. [231]/1997	50	Diagnostic and operative laparoscopy	Needle: 3.8 Open: 1.8	Needle: 0/25 Open: 0/25	Open technique is faster
Byron et al. [39]/1993	252	Diagnostic and operative laparoscopy	Needle: 5.9 Open: 2.2	Needle: 19/141 Open: 4/111	Open technique is safer and faster
Nezhat et al. [219]/1991	200	Diagnostic and operative laparoscopy	Not mentioned	Needle: 22/100 Open: 3/100	Open technique has fewer complications
Borgatta et al. [30]/1990	212	Laparoscopic tubal sterilization	Needle: 9.6 Open: 7.5	Needle: 7/110 Open: 4/102	Open technique is safer and faster

Table 3. Randomized clinical trials comparing low- and high-pressure pneumoperitoneum^a

Reference/year	No. of patients ASA	Pressures compared	Results	Conclusions
Wallace et al. [308]/1997	40/ASA I–II	7.5 vs 15 mmHg CO ₂	CI↓, MAP↑, HR↓, end-tidal CO ₂ ↑, pain scores↓	Cardiac changes in both groups similar; postop pain in low-pressure group reduced
Pier et al. [236]/1994	33/ASA I–II	8 vs 15 vs 19 mmHg CO ₂	No differences in pain, analgesic use, FEV ₁ , or VC	Pressure has little effect on pain
Dexter et al. [63]/1999	20/ASA I–II	7 vs 15 mmHg CO ₂	MAP↑, HR↑, SV↓, CO↓	High pressure reduces SV and CO more than low PP

CO, Cardiac output; HR, heart rate; MAP, mean arterial pressure

^a All trials were performed on laparoscopic cholecystectomy

procedures, although one study did not confirm this result (1b [112]). By comparisons carbon dioxide insufflation with gasless laparoscopy, similar courses of stress response parameters were found (1b [158, 162]), but conflicting data exist (1b [126]). Since all these studies compared laparoscopic and open surgery, the immunological effects of the pneumoperitoneum and the surgical procedure overlap each other, precluding the quantification of any specific effects.

The influence of the specifics of the pneumoperitoneum (e.g., IAP, gas, and warming and humidified surrounding) on immunological function has only partly been studied in experimental settings. Helium seems to preserve cell-mediated intraperitoneal immunity better than CO₂ (5 [47, 219]) and causes a less pronounced cytokine response and bacterial translocation (5 [194]). In clinical trials, postoperative intraperitoneal cytokine response after warming the insufflation gas was attenuated (1b [244]). Another study suggested a similar stress response (IL-6, CRP, neutrophil elastase, and white cell count) after pneumoperitoneum or abdominal wall lifting (1b [221]). It is questionable whether the specifics of the pneumoperitoneum have clinically relevant effects or even benefits on postoperative immunological function and outcome (e.g., less pain, fatigue, and complica-

tions). Thus, additional clinical trials with adequate end points and sample sizes have to be performed to confirm the hypothesis of better preservation of the immune function by minimally invasive surgery.

Peritonitis

Presupposing appropriate perioperative measures (e.g., adequate preoperative volume loading) and hemodynamic stability, there are no contraindications to create a pneumoperitoneum when laparoscopic surgery is applicable in cases of peritonitis (grade B). The results from animal studies on the influence of pneumoperitoneum bacteremia and endotoxemia are controversial.

In experimental studies, a pneumoperitoneum seems to increase the risk of bacteremia and endotoxemia [23–25, 77, 101, 214]. Other animal studies demonstrated that the systemic inflammation is higher after laparotomy than after laparoscopy, causing a transient decrease in immunologic defense and possibly leading to sepsis (5 [131, 180]).

With regard to the specifics of a pneumoperitoneum, any increase in IAP seems to further promote bacteremia (2b [77]), but data are inconsistent. The used gas

Table 4. Pneumoperitoneum and hypothermia; randomized clinical trials

References/years	No. of patients operations	Treatments	Temperature measurement	Pathophysiological results	Clinical results
Dietterle et al. [323]/1998	100 vs 100 operative or diagnostic pelviscopic procedures	Body vs room temperature, pressure and humidifying not mentioned	None	None	Gas warming lowers the intensity of diaphragm and shoulder pain and reduces the use of analgesics
Saad et al. [254]/2000	10 vs 10 lap. CCE	37 vs 21°C. CO ₂ IAP 15 mm Hg, humidifying not mentioned	Esophageal thermotip catheter	No differences in body and intraabdominal temperatures and pain scores	Gas warming has no clinically relevant effect
Slim et al. [284]/1999	Double-blind 49 warm vs 51 cold gas, different upper abdominal lap. procedures	37 vs 21°C, CO ₂ , IAP 14 mmHg humidifying not mentioned, 20°RT	Subdiaphragmatic thermometric probe	Subdiaphragmatic temperature equal VAS score for shoulder tip pain higher in the warm group	Gas warming increases postoperative pain (VAS)
Mouton et al. [209]/1999	20 vs 20, lap. CCE	34–37 vs 21–25°C	Esophagus	No difference in core temperature; pain score less in humidified group	Humidified heated gas reduces pain but preserves no heat loss
Nelskylä et al. [215]/1999	18 vs 19 women, lap, HE End point: heart rate variability	37 vs 24°C, CO ₂ , IAP 12–14 mmHg, humidifying not mentioned	Tympanic and nasopharyngeal	Body core temperature decreases more in the warming group	Heating of insufflation gas does not prevent decrease of body temperature
Puttick et al. [243]/1999	15 vs 15, ASA I–II, lap CCE	37 vs 21°C, CO ₂ Mean duration of surgery 32 min	Esophagus	Body core temperature decreases more in room temperature group	Higher cytokine levels in room temperature group; pain scores and consumption not different
Bäcklund et al. [111]/1998	13 vs 13 prolonged lap. procedures > 90 min	37 vs 21°C, CO ₂ , IAP 11–15 mmHg, humidifying not mentioned	Swan–Ganz catheter	Core temperature and urine output higher in warm PP	Warm insufflation increases urine output
Ott et al. [226]/1998	Double-blinded multi-center (7) study 72 women	36°C and humidified vs 23°C, CO ₂ IAP?	Endotracheal	Warm insufflation: less intraoperative hypothermia post-operative stay and pain	Gas warming reduces lap. induced hypothermia
Korell et al. [162]/1996	50 vs 53 women, div. laparoscopic procedures	Heated CO ₂ (30–32°C) vs normal CO ₂ (23–24°C)	Flow therm	VAS scores reduced for shoulder and subdiaphragmatic pain	Warm CO ₂ reduces postoperative pain
Semm et al. [277]/1994	30 vs 30 lap. pelviscopy	37°C PP vs 21°C PP, CO ₂ IAP 12 mmHg, humidifying not mentioned	Intraabdominal and rectal probe	37°C group shoulder tip pain; pain medication and incidence of tachycardia reduced	Warm insufflation reduces shoulder tip pain; pain medication

IAP, intraabdominal pressure

seems to play only a minor role (5 [105]). A clinical RCT found no difference in the acute phase response and endotoxemia between laparoscopic and conventional gastric surgery in cases of peritonitis (1b [173]). Furthermore, laparoscopic compared to conventional cholecystectomy for acute and gangrenous cholecystitis does not increase the mortality rate (1b [153]), and the morbidity rate seems to be even lower after laparoscopy (1b [153, 182]). Two small conflicting RCTs assessed bacteremia during appendectomy and found 0/11 versus 6/12 and 5/14 versus 5/13 positive blood cultures after open and laparoscopic access, respectively (1b [222, 279]). The hypothesis that in cases of peritonitis laparoscopy leads to a lesser depression of the systemic immune response with better postoperative outcome is unproven.

In conclusion, the decision to perform a laparoscopic procedure in case of peritonitis depends on the extent of peritonitis, the onset of disease, and the general clinical state of the patient. No clinical trials have found any contraindication to perform laparoscopy in case of beginning peritonitis (e.g., perforated appendicitis).

Risk of tumor spreading

There is no strong clinical evidence (except case reports) that pneumoperitoneum in patients with intraabdominal malignant disease increases the risk of tumor spread (grade D). The panel considers that there is no reason to contraindicate pneumoperitoneum in these patients, given the fact that an appropriate operative technique is used

Table 5. Randomized clinical trials comparing gasless to low- or high-pressure pneumoperitoneum

Reference/year	No. of patients, ASA	Pressures compared	Results in experimental group	Conclusions
Lubkan et al. [184]/2000	30, single blind	Laparolift vs 15 mmHg	VAS score for visualization less in gasless patients	Conventional PP provides better view
Ogihara et al. [223]/1999	12, ASA I–II	Gasless vs 13 mmHg CO ₂ (Trendelenburg position 15–20° in both groups)	CO ₂ group: pulmonary compliance↓, epinephrine↑, norepinephrine↑, dopamine↑, ADH↑, urine output↓	Gasless: lesser hormonal stress responses; better pulmonary function; higher urine output
Schulze et al. [271]/1999	17	Gasless vs 12 mmHg CO ₂ (thPDA in both groups)	CO ₂ group Blood flow↓, HR↑, MAP↑, CVP↑,	No clinically relevant differences; CO ₂ group less pain and more fatigue
Vezakis et al. [304]/1999	36, ASA I–II	Gasless vs 8 mmHg CO ₂	No changes in postop pain and analgetic consumption	Shoulder pain more frequent
Cravello et al. [53]/1999	103	Gasless vs CO ₂ PP (IAP unknown) (8 conversions)	No differences in complication pain medication hospital stay	Gasless technique needs further evaluation
Guido et al. [102]/1999	54	Gasless vs 15 mmHg CO ₂	No differences in shoulder, pelvic, and periumbilical pain	Similar pain scores compared to conventional PP
Meijer et al. [200]/1997	20, ASA I–II	Gasless (AWL) + 5 mmHg vs 15 mmHg CO ₂	Gasless surgery lasted longer; CO, RR, and HR equal in both groups	AWL is not recommended for laparoscopic cholecystectomy, view impaired
Koivusalo et al. [156]/1997	30, ASA I–II	IAP 12–13 mmHg CO ₂ vs gasless	CO ₂ group: MAP↑, pulmonary compliance↓, urine output↓, U-NAG↑, intramucosal pH↓	Gasless: more stable in hemodynamics; protects renal and splanchnic ischemia
Koivusalo et al. [155]/1997	25, ASA I–II	IAP 12–15 mmHg CO ₂ vs gasless AWL	Drowsiness shorter	Avoiding CO ₂ reduces drowsiness
Johnson and Sibert [136]/1997	18	Gasless vs CO ₂ PP (IAP not mentioned)	Increased technical difficulty—poor visualization	CO ₂ PP is preferable for routine LTC
Casati et al. [42]/1997	20	Gasless vs 12 mmHg CO ₂	Better pulmonary compliance: oxygenation unchanged	Better lung compliance
Goldberg and Maurser [96]/1997	57	Gasless (laparolift) vs 15 mmHg CO ₂ ; 9/28 converted because of inadequate exposure	Technically difficult; no differences in cardiopulmonary parameters and pain scores	No clinical benefit
Koivusalo et al. [154, 157]/1996	26, ASA I–II	IAP 12–15 mmHg vs laparolift	Maddox–Wing deviation higher in conventional PP group gasless: plasma renin activity↓, diuresis higher	Less right shoulder pain, nausea, and vomiting; smaller neuroendocrine responses; better renal function
Lindgren et al. [178]/1995	25, ASA I–II	PP (IAP 12–15 mmHg + CO ₂) vs AWL	CO ₂ group MAP↑, HR↑, pulmonary compliance↓	MAP lower; postoperative nausea, vomiting, and right shoulder pain less often

AWL, abdominal wall lifting; CO, cardiac output; HR, heart rate; MAP, mean arterial pressure

(grade C). The type of insufflation gas seems to affect intraabdominal tumor growth, whereas intraabdominal pressure is of little importance (grade D). Due to the low level of evidence, patients undergoing laparoscopic surgery for malignant disease should be included in randomized controlled trials or at least in quality registries.

Several animal studies have been conducted to evaluate the pathogenesis of portsite metastasis in laparoscopic surgery, but the experimental models and tumor cell techniques vary considerably (5 [32, 33, 132, 134, 151, 219]). Port-site recurrence is common in small animal models after inoculation of high numbers of tumor cells and more pronounced after capnoperitoneum compared to laparotomy or gasless laparoscopy (5 [132, 134, 219]). IAP has little influence on intraperitoneal tumor growth or the incidence of port-site metastasis, whereas insufflation with helium may decrease subcutaneous tumor growth (5 [132, 134, 219]). In contrast to these findings, intraperitoneal tumor growth is stimulated more by laparotomy than by laparoscopy, gasless laparoscopy, or anesthesia alone without any operative procedure (5[130]).

Port dislocations should be avoided and ports should be irrigated intraperitoneally before they are retracted from the abdominal cavity. Before the tumor is extracted, the incision has to be protected against direct tumor cell contamination. The risk of tumor cell dissemination may be reduced by intraabdominal instillation of cytotoxic solutions at the end of operation (5 [34]).

Prospective clinical trials failed to show a higher incidence of free intraperitoneal tumor cells (5 [37]) or recurrence in the skin incision (2a [304], 5 [37]) for laparoscopic compared to conventional surgery. A systematic review of clinical trials found no significant differences in overall survival, disease-free survival, cancer-related death, locoregional tumor recurrences, port-site metastasis, or distant metastasis in patients undergoing laparoscopic or conventional colorectal resections (2a [304]). Perioperatively, mobilization of neoplastic cells occurs frequently in patients with colorectal cancer, but the surgical approach does not seem to be a determining factor (1b [18]). Randomized trials with low quality found no wound or port-site metastasis

Table 6. Cross-tabulation of current research on the effects of technical modifications of laparoscopy on pathophysiologic and medical outcomes

	Open-access technique	Smaller trocars (3.5 mm)	Warmed insufflation gas	Helium, argon, or NO ₂	Low-pressure laparoscopy	Gasless laparoscopy	Intraperitoneal anaesthetics
Pathophysioloical effects							
Circulatory	(0)	(0)	++ [277]	0 [28]	++ [63, 158, 236, 308]	++ [96, 156, 158, 178, 200, 220, 223]	(0)
Pulmonary	(0)	(0)	?	0 [28]	+ [236]	++ [155, 158, 200, 220, 223]	?
Renal/Hepatic/Intestinal	(0)	(0)	++ [11]		+	++ [156, 158, 223]	(0)
Immunological	(0)	?	++ [243]	(0)	(0)	+/0 [220]	?
Hormonal	(0)	?	?	(0)	(0)	++ [156, 223]	?
Body core temperature	(0)	(0)	+/0 [11, 215, 226, 243, 254]	?	(0)	+	(0)
Technical effects	++ [262]	- [276]	(0)	(0)	-/0 [63, 158, 308]	- [96, 136, 184, 200]	(0)
Clinical effects							
Intraoperative surgical incidents	++ + [30, 39, 50, 103, -219, -231]	[276]	+	(0)	(0)	?	(0)
Heart and lung complications	(0)	+ [276]	?	(0)	-10	+/0	?
Kidney and liver complications	(0)	(0)	?	(0)	-10	+/0	?
Pain	(0)	++ [22, 35, 276]	+/0 [162, 243, 254, 277, 284]	0 [236]	++ [258, 308]	+/0 [53, 102, 154, 178, 271, 304]	+++ [3, 40, 49, 55, 69, 70, 93, 211, 214, 227, 228, 293, 297, 309]
Drowsiness and fatigue	(0)	0 [276]	?	(0)	(0)	++ [155, 178, 271]	?
Cosmetic results	?	++ [276]	(0)	(0)	(0)	(0)	(0)
Incisional hernia	?	++	(0)	(0)	(0)	(0)	(0)
Adhesions	(0)	+	(0)	(0)	+	(0)	(0)
Infections	(0)	?	(0)	(0)	(0)	(0)	(0)

+++ , strong RCT evidence in favour of intervention; ++ , some RCT evidence in favour of intervention; +/0, conflicting RCT evidence in favour of intervention; + , non-RCT evidence in favour of intervention; 0, some RCT evidence for no effect of intervention; (0), non-RCT evidence for no effect of intervention; - , non-RCT evidence against intervention; -/0, conflicting RCT evidence against intervention; - some RCT evidence against intervention; — , strong RCT evidence against intervention; ? , no valid research evidence available

in 91 patients during a mean follow-up of 21.4 months and in 43 patients after long-term 5-year follow-up (2b [57, 169]). Adequately powered RCTs on laparoscopic and conventional resections of colorectal carcinoma are missing, but such trials are currently being performed in Europe, the United States, and Australia. Results of these trials will be available in 2004–2006.

Establishing the pneumoperitoneum

Creation of a pneumoperitoneum

For severe complications (vessel perforation) it is impossible to prove a difference between closed- and open-access technique in RCTs; therefore, large outcome studies should be considered. In the RCTs, the rate of major and minor complications is surprisingly high, which may be due to the definition of a complication or surgical learning

curve. Insertion of the first trocar with the open technique is faster as compared to the Veress needle (grade A). The randomized controlled trials comparing closed (Veress plus trocar) versus open approach have inadequate sample sizes to find a difference in serious complications. In large outcome studies there were less complications in the closed group (grade B). Although RCTs found the open approach faster and associated with a lower incidence of minor complications (grade A), the panel cannot favor the use of either access technique. However, the use of either technique may have advantages in specific patient subgroups (grade B).

Among the various techniques for achieving a pneumoperitoneum and introducing the first trocar, two common methods are usually performed. The first, so-called closed technique requires the Veress needle, which is inserted in the abdominal cavity for CO₂ insufflation followed by blind introduction of the first trocar. The second, so-called open technique was first described by

Hasson [110]. This technique begins with a small incision at the umbilical site and subsequently all layers of the abdominal wall are incised. The first trocar is then inserted under direct vision followed by gas insufflation.

The morbidity associated with the establishment of the pneumoperitoneum and the insertion of the first trocar is estimated to be less than 1% (4 [29, 109, 264]), but the true incidence of visceral and vascular injury for both techniques is unknown. However, major vascular injuries occur most often with the Veress needle (2c [44, 236]). Several RCTs found that the open technique on average causes less complications and is cheaper and faster than the Veress needle technique (1b [30, 39, 50, 104, 220, 232]) (Table 2). However, one study on the access technique for percutaneous diagnostic peritoneal lavage in blunt trauma patients showed that the Veress needle technique was faster compared to the open technique (1b [263]). A recent three-armed RCT found it easier to establish the pneumoperitoneum with a new access device (TrocDoc) than with the open technique or the Veress needle (1b [14]). The choice between reusable and single-use instruments was outside our scope. In specific patient subgroups, the access technique has to be chosen according to the patients characteristics (e.g., pregnancy, obesity, and trauma).

Gas embolism and its prevention

Clinically relevant gas embolism is very rare, but if it occurs, it may be a fatal complication (grade C). The true incidence of clinically inapparent gas embolism is not known. Most described cases of gas embolism have been caused by accidental vessel puncture with a Veress needle at the induction of pneumoperitoneum. Low intraabdominal pressure, low insufflation rates, as well as careful surgical technique may reduce the incidence of gas embolism (grade D). A sudden decrease in end-tidal CO₂ concentration and blood pressure during abdominal insufflation should be considered a sign of gas embolism (grade C). Due to the low incidence of clinically relevant gas embolisms, advanced invasive monitoring (transesophageal Doppler sonography) cannot be recommended for clinical routine (grade B).

The incidence of gas embolism during pneumoperitoneum is estimated to be less than 0.6% (2 [282], 4 [122, 144]). Many case reports have detailed fatal or near-fatal coronary, cerebral, or other gas embolism (4 [102, 152, 172, 231, 238]). In more than 60% of cases, gas embolism occurred during the creation of a pneumoperitoneum. The usual cause leading to gas embolism was the accidental displacement of a needle or trocar into a blood vessel. Similarly, any injury to the veins of parenchymal organs can result in direct gas flow into systemic circulation. CO₂ bubbles are capable of reaching the right heart (2b–5 [61, 66, 79, 267]). This is best detectable when patients are studied with transesophageal echocardiography (2b–5 [61, 66]). Transcranial Doppler has shown that CO₂ bubbles may even reach the cerebral circulation (4 [267]). Furthermore, gas emboli are able to escape from venous to arterial circulation through pul-

monary arteriovenous shunts (5 [306]) or an open Foramen ovale (4 [190]).

Experimental animal studies have induced gas embolism by infusing air directly into a vein or by lacerating a large intraabdominal vein during a pneumoperitoneum (5 [66, 145, 147]). Increased IAP of more than 20 mmHg in connection with an insoluble gas (helium or argon) enhanced the risk of gas embolism during pneumoperitoneum (5 [146, 148, 251]), suggesting that caution should be exerted when laparoscopic surgery is performed close to large veins (5 [66]). Furthermore, the use of nitrous oxide for anesthesia may increase the risk of developing gas embolism during laparoscopy (4 [200, 242], 5 [147]).

In clinical practice, there are few technical options available to reduce the risk of gas embolism. It is therefore very important that especially the surgeon who creates the pneumoperitoneum be experienced in laparoscopic access techniques. It can be assumed that blunt trocars reduce the risk of accidental vessel puncture (1b [14]).

The most sensitive method to detect gas embolism is transesophageal Doppler monitoring (TEE) (2b [283, 316]). Simple measures to detect clinically relevant gas embolism are electrocardiogram (ECG) and EtCO₂ monitoring, which have low costs and require low personal effort. During surgery, decreasing EtCO₂ values of more than 3 mmHg could be related with gas embolism and should be clarified immediately (4 [52], 5 [147]). In case of injury of larger veins during abdominal insufflation, ECG and EtCO₂ should be closely monitored, especially when gases with low solubility are used. Because of the low incidence of gas embolism, special perioperative monitoring (e.g., TEE) is not indicated.

Choice of insufflation pressure

The panel recommends use of the lowest IAP allowing adequate exposure of the operative field rather than using a routine pressure (grade B). An IAP lower than 14 mmHg is considered safe in a healthy patient (grade A). Abdominal wall-lifting devices have no clinically relevant advantages compared to low-pressure (5–7 mmHg) pneumoperitoneum (grade B).

Normal and low laparoscopic insufflation pressure are defined as 12–15 and 5–7 mmHg, respectively. It is important to differentiate between the pressure at induction of the pneumoperitoneum and that during the operation. Initially, the IAP might be increased up to 15 mmHg to reduce the risk of trocar injuries. As already stated, IAP affects the physiology of heart, lung, and circulation. In order to attenuate these possible side effects of high IAP, the intravascular volume should be adequately filled preoperatively (1b [159]) and the insufflation pressure should be selected according to the planned laparoscopic procedure and the patient characteristics. In ASA I and II patients, a low-pressure pneumoperitoneum reduces adverse effects on physiology without compromising laparoscopic feasibility (1b [63, 237, 309]) (Table 3). It remains questionable

whether these physiologic changes are associated with clinically relevant side effects.

In older and compromised patients (ASA III and IV), the effects of a high vs low IAP have only been studied in nonrandomized clinical trials (2b [64, 83, 111], 4 [257]). In these studies, an elevated IAP (12–15 mmHg) showed considerable cardiac alterations. With the use of invasive monitoring, adequate volume loading, and vasoactive drug, it was possible to keep the hemodynamic and cardiac function stable. Therefore, in ASA III and IV patients, gasless or low-pressure laparoscopy could be alternatives, which should be further tested.

Warming and humidifying of insufflation gas

Warming and humidifying insufflation gas is intended to decrease heat loss. However, compared to external heating devices, the clinical effects of warmed, humidified insufflation gas are minor (grade B). Data on its influence on postoperative pain are contradictory (grade A).

Perioperative hypothermia is related to increased catecholamine and cortisol levels leading to peripheral vasoconstriction and higher arterial blood pressures (2b [86]). Maintaining normothermia generally decreases postoperative cardiovascular morbidity (1b [84, 85]).

General and regional anesthesia essentially determine body core temperature by downregulation of the internal temperature level. Once vasoconstriction has occurred, application of warming systems is less effective in compensating heat loss (1b [245]). Therefore, forced-air warmer systems should be applied before heat loss occurs. In contrast, warming and humidifying of the insufflation gas is less important than application of external warming devices before and during anesthesia.

Warming of the insufflation gas reduces postoperative intraperitoneal cytokine response (1b [243]) and reduces postoperative hospital stay (1b [226]) and pain (1b [226, 277, 323]) (Table 4). In contrast, a double-blind RCT found an increase in shoulder tip pain after warming the insufflation gas (1b [284]). Other groups found no clinically relevant effects of warming the insufflation gas (1b [198, 215, 254, 311]). Additional humidifying of warmed insufflation gas seems to reduce postoperative pain but has no heat-preserving effect in brief laparoscopic procedures (1b [209]). Since most of the studies have small sample sizes with possible type II error, no firm conclusions can be drawn. Given their possible small effects, the costs of these devices have also to be considered.

Abdominal wall-lifting devices

Abdominal wall lifting as compared to capnoperitoneum results in less impairment of hemodynamic, pulmonary, and renal function (grade A). In ASA and I and II patients, the magnitude of these benefits is too small to recommend abdominal wall lifting (grade D). In patients with limited cardiac, pulmonary, or renal function, abdominal wall lifting combined with low-pressure pneumo-

peritoneum might be an alternative (grade C). Nevertheless, surgical handling and operative view were impaired in most surgical procedures (grade A).

Gasless laparoscopy has been developed to avoid the pathophysiological side effects of elevated IAP and CO₂ insufflation, especially in patients with comorbidities (ASA III and IV). However, most RCTs on gasless laparoscopy vs pneumoperitoneum have been performed in healthy ASA I and II patients (Tables 5 and 6). In these patients, gasless laparoscopy results in a more stable hemodynamic and pulmonary function (1b [155, 220, 223]), a concomitant increase in urine output (1b [156, 223]), reduced hormonal stress responses (1b [156, 223]), less postoperative pain (1b [131, 153]), and less drowsiness (1b [155, 178]). Contrarily, other RCTs found no differences in postoperative pain (1b [102]) and cardiorespiratory functions (1b [200]). Many surgeons encountered technical difficulties due to inadequate visualization (1b [136, 184, 200]). This led to high conversion rates in these trials, one of which was even terminated prematurely [136]. Although gasless laparoscopy may have hemodynamic and cardiovascular advantages in ASA III and IV patients, clinical trials in this group of patients have not been performed. Since gasless laparoscopy also requires excellent surgical expertise, its use should be restricted to certain subgroups of surgeons and patients.

Size of access devices

Smaller access devices (≤5 mm) in laparoscopy are only feasible in selected group of patients. The use of 2 to 5-mm instead of 5 to 10-mm access devices improves cosmetic result and postoperative pain marginally in laparoscopic cholecystectomy (grade A).

Although it has been assumed that smaller access devices may markedly improve the patients outcome of laparoscopic surgery, this has not been shown in valid RCTs (1b [22]). Merely modest advantages have been reported concerning a better cosmetic result (1b [276]) and less postoperative pain (1b [22, 35, 46, 276], 4 [192]) after laparoscopic cholecystectomy. Postoperative pulmonary function and fatigue were unchanged (1b [276]). Other clinical trials found a shorter convalescence by using smaller access devices in laparoscopic procedures (4 [192]). The incidence of postlaparoscopic incisional hernia is less than 1% (4 [165, 169]). Whether smaller access devices prevent incisional hernia has not been clarified (4 [165]). To prove a difference would require a large sample size and an extensive postoperative observation period. Currently, the general use of smaller trocars cannot be recommended due to difficulties in handling and reduced optical quality, especially when using smaller laparoscopes (1b [22, 276], 4 [168]). Recently published RCTs reported a reduction in postlaparoscopic pain when a radially expanding access device was compared to the conventional cutting trocar (1b [19, 80, 170, 318]). No data are available on other clinical effects.

Perioperative aspects

Adhesions

Some laparoscopic procedures may cause less postoperative adhesions compared to their conventional counterparts (grade B). However, the specifics of a pneumoperitoneum (gas, pressure, temperature, and humidity) seem to have no major effect on the development of postsurgical adhesions (grade D).

Two RCTs found less postsurgical adhesions after laparoscopic compared to conventional surgery (2b [61, 62, 185]), but these studies have methodological flaws (small sample size, unclear allocation concealment, no intention-to-treat analysis, and losses to follow-up). Furthermore, since postsurgical adhesions are usually assessed by means of different scoring systems, it is difficult to compare the results of the trials in between or to rule out observer bias in these unblinded trials.

Pathophysiologically, a reduced peritoneal fibrinolytic activity seems to be the main cause for postsurgical adhesions (4 [116]). Experimental studies indicate that adhesion rates also depend on intraabdominal pressure (5 [317]) and the type of gas used (5 [132, 213]). However, one clinical RCT found no difference in peritoneal fibrinolytic activity in elective laparoscopic compared to conventional colorectal resections (1b [216]). It seems that the specifics of a pneumoperitoneum do not influence generally the peritoneal fibrinolytic activity and the development of postoperative adhesions. Therefore, avoiding local peritoneal damage seems to be the most significant factor to prevent postsurgical adhesions.

Pain, nausea, and vomiting

Pain after laparoscopic surgery is multifactorial and should be treated with a multimodal approach (grade A). Shape and size of access devices have to be considered (grade A). Low-pressure pneumoperitoneum, heated and humidified insufflation gas, incisional and intraperitoneal instillation of local anesthetics, intraperitoneal instillation of saline, and removal of residual gas all reduce postlaparoscopic pain (grade B). Inconclusive data and small effect sizes of singular approaches make it difficult to recommend these treatments in general (grade D). No evidence exists that the specifics of a pneumoperitoneum have any effect on postoperative nausea and vomiting.

Although pain after laparoscopic surgery is less severe and of shorter duration than that after open surgery, it still causes considerable discomfort and increased stress response. The etiology of postlaparoscopic pain can be classified into at least three aspects: visceral, incisional, and shoulder pain [21, 140, 300]. Although visceral pain may also depend on the extent of intraabdominal surgery, incisional pain is related to the number and size of access devices and also to the technique of incision closure and drainage. The origin of shoulder pain is only partly understood, but it is commonly assumed that the continual stretching of

the peritoneum during and after the pneumoperitoneum is responsible. Clinically, incisional and deep abdominal pain dominate over shoulder pain. However, shoulder pain is specific for laparoscopic surgery. After different abdominal laparoscopic procedures, shoulder pain was noted in 30–50% of cases, which is significantly higher than after the corresponding open procedures (1b [43, 55, 174, 297]). It was suggested that shoulder tip pain is caused mechanically by stretching the diaphragmatic ligaments (1b [308]). The hypothesis of a chemical effect of the pneumoperitoneum with a decrease in intraperitoneal pH could not be verified [233].

The incidence of postoperative nausea and vomiting after laparoscopic procedures ranges from 10 to 60% [81, 201, 312]. After laparoscopic cholecystectomy, nausea and vomiting persisted up to 14 days in some patients [296]. The pathogenesis of postoperative nausea and vomiting is multifactorial, depending on anesthesia, surgery, gender, and perioperative administration of opioids. Several RCTs examined the influence of antiemetics and analgesics on postoperative nausea and vomiting, but this was beyond the scope of this guideline.

Within the past few years, various modifications of the pneumoperitoneum have been developed and clinically tested in order to reduce peritoneal pain after laparoscopic surgery [21, 310]. Here, we focus on those interventions that are directly related to the pneumoperitoneum, thus excluding oral, intravenous, or epidural drug administration and other nonlocal treatments. The intensity of postoperative pain varies largely among different cultures, settings, and individuals.

The following interventions were all shown in RCTs to effectively reduce pain after laparoscopy:

- Reducing IAP (1b [178, 236, 299, 308])
- Using other insufflation gases, such as N₂O, helium, or argon (1b [2, 180, 206, 236, 280]),
- Lowering the insufflation rate (1b [16])
- Warming and humidifying the insufflation gas (1b [162, 209, 210, 226, 254])
- Removal of residual intraabdominal gas at the end of operation (1b [86, 137, 298], 2b [4, 128], 4 [4, 128])
- Intraperitoneal instillation of fluids (1b [233])
- Intraperitoneal instillation of anesthetics (1b [3, 40, 49, 55, 69, 70, 93, 211, 214, 227, 228, 293, 297, 309])
- Reducing the size of trocars (1b [22, 35, 97, 166])
- Injecting anesthetics into the trocar sites (1b [3, 21, 257, 301])
- Omitting drains (which is beyond the scope of this recommendation, since it depends on the type of operation)

The intraperitoneal instillation of anesthetics is well studied. Most RCTs found a significant decrease in postlaparoscopic pain, including shoulder tip pain (1b [3, 40, 49, 55, 69, 70, 93, 211, 214, 227, 228, 293, 297, 309]), whereas other trials found no effect (1b [21, 87, 140, 244, 264, 270]). Since there is also evidence that postlaparoscopic instillation of normal saline or Ringers lactate reduces pain (1b [233]), it is important to distinguish between trials that used placebo controls from those that did not.

Humidifying the insufflation gas reduced postoperative pain in one trial (1b [211]) but increased it in another (1b [284]). After gasless laparoscopy, one double-blind RCT showed that shoulder tip pain (as primary end point) was more frequent than after conventional pneumoperitoneum (1b [304]), a second RCT with a smaller sample size reported the contrary (1b [154]), and a third found no difference (1b [101]).

On the basis of these contradictory results, the panel is not able to favor one treatment option over another. For the multifactorial pathogenesis of postlaparoscopic pain, we assume that a combined therapeutic approach is most effective ([20, 201]). Surgical awareness of this significant patient problem needs to be improved.

Pregnancy

Presupposing obstetrical consultation, laparoscopic procedures during pregnancy should be performed in the second trimester if possible (grade C). Perioperatively, maternal end-tidal CO₂ concentration and arterial blood gases must be monitored to control maternal hyperventilation and to prevent fetal acidosis (grade C). For the establishment of the pneumoperitoneum the open technique should be preferred (grade C). During laparoscopy intraabdominal pressure should be kept as low as possible and body positioning should be considered in order to avoid inferior vena cava compression by the uterus (grade C). Furthermore, pneumatic compression devices are recommended (grade D).

Surgery during pregnancy always carries an increased risk of fetal loss. Therefore, the indication for surgical intervention during pregnancy is generally limited to urgent situations such as acute appendicitis [268] or acute cholecystitis [99, 287]. The incidence of acute appendicitis and acute cholecystitis during pregnancy is similar to that of nongravid females and is estimated to be less than 0.1% (4 [195, 248]). The treatment of acute cholecystitis in gravid women should consider effective nonsurgical therapeutic options (4 [292]). Today, pregnancy should not be seen as an absolute but a relative contraindication for laparoscopic procedures. Because of increased risk for postoperative abortion in the first trimester and hindrance of operation due to the enlarged uterus, surgery during pregnancy should be performed during the second trimester (4 [190, 286]). During pregnancy laparoscopic compared to conventional surgery is preferred because of possibly less fetal impairment due to less postoperative analgetic requirements (4 [176]) and less postoperative maternal respiratory depression (4 [56]). However, increased intraabdominal pressure may decrease maternal respiratory compliance (5 [9, 58]), uterine blood flow (5 [58]), or preterm labor (5 [58, 96]). Furthermore, the use of carbon dioxide seems to increase fetal acidosis (5 [54, 59, 120]), to enhance the risk for fetal loss (4 [8]), and may lead to detrimental side effects if hyperventilation fails (5 [54]). Most of these concerns are based on experimental studies and case reports and should be confirmed by randomized controlled trials. Due to the low incidence of surgical interventions during preg-

nancy, these studies have to be performed as multicenter trials.

Intracranial pressure

Increased IAP and head-down position increase intracranial pressure (ICP) (grade A). Therefore, elevated IAP, head-down position, and hypoventilation should be avoided (grade D). In patients with head injury or neurological disorders, perioperative monitoring of ICP should be considered (grade C). Gasless laparoscopy might be an alternative to prevent ICP peaks (grade D).

During pneumoperitoneum, IAP and head-down position increase ICP (5 [75, 117, 142, 207, 252], 4 [123]), enhance cerebral blood flow velocity (4 [1]), and diminish cerebrospinal fluid absorption (5 [106]). Elevated ICP values during laparoscopic surgery return to baseline after desufflation (5 [75]). There is no evidence that elevated ICP during pneumoperitoneum is clinically relevant.

Pathophysiological studies suggested that an increased intraabdominal pressure hinders venous drainage of the lumbar venous plexus followed by a decline in cerebrospinal fluid absorption during abdominal CO₂ insufflation (5 [105, 106]). Furthermore, it was hypothesized that this mechanical effect leads to an increase in ICP and a central nervous system response causing systemic hypertension (5 [15, 251]). However, the exact pathophysiology of increasing ICP during pneumoperitoneum remains unclear. Experimental and clinical studies showed that hemodynamic changes are directly related to the increase in ICP (4 [89, 123], 5 [142, 251]). Therefore, in patients with severe head injuries or conditions associated with elevated ICP, intraabdominal pressure should be as low as possible, sudden IAP peaks should be avoided, and intraoperative ICP monitoring should be considered (4 [127]). Furthermore, gasless laparoscopy could be an option to avoid the effects of IAP on ICP (5 [75, 117]).

The use of carbon dioxide as insufflation gas leads to hypercarbia and acidosis, which possibly influence the intracerebral circulation by vascular autoregulation. CO₂ increases ICP more than do helium and nitric oxide (5 [267]). Hypoventilation and hypercarbia increase ICP compared to hyperventilation and hypocarbia, but during acute elevations of ICP hyperventilation did not decrease ICP effectively (5 [251]). The insufflation gas has fewer effects on ICP than on IAP (4 [74], 5 [60]).

Abdominal trauma

There are no prospective studies evaluating the specifics of a pneumoperitoneum (type of gas, IAP, and temperature) in patients with blunt or penetrating abdominal trauma (grade D).

Laparoscopy is used as a diagnostic tool in hemodynamically stable patients after blunt or penetrating trauma in order to detect those injuries that require laparotomy or laparoscopic repair (2b [71, 77, 285]). In rare cases of penetrating trauma, the establishment of a

pneumoperitoneum led to an insufflation of injured organs or cavities ([119]). Nevertheless, the panel agrees that there is no reason to contraindicate pneumoperitoneum in stable trauma patients.

The use of different intraabdominal pressures, different types of gas, or even gasless laparoscopy has not been evaluated in patients with blunt or penetrating abdominal trauma. Thus, no recommendations are reasonably justifiable. However, one clinical RCT tested which access technique is faster and safer, and found advantages for the closed technique (1b [262]), thus refuting data from nontrauma surgery.

Discussion

After a 2-year break, the E.A.E.S has continued its guideline activities, now on an even more evidence-based level and with much more advanced preparation than in the past. We believe that the result of this endeavor can be considered to be a milestone in the society's responsibility of being a bridgetender between primary research and clinical practice and vice versa.

We hope that the reader understands the importance of guideline methodology. In a European survey 2 years ago, many members complained that the E.A.E.S consensus panels had always been consisting of the same clique of people. The panel for this guideline still contains many well-known names from the E.A.E.S simply because the number of experts in endoscopic surgery is limited, as are resources for guideline development. Wherever interdisciplinary coworking was necessary, experts from other fields were invited to join the panel, although this guideline could have received further benefit from the input of a pediatric surgeon.

The scope of this guideline is broad since the pneumoperitoneum is the key issue in laparoscopic surgery. However, it is impossible for a guideline to answer all relevant points in detail or to discuss the role of the pneumoperitoneum separately for every disease entity. The panel tried to formulate the statements as concise as possible. However, for those issues, for which no strong evidence was found, it was often impossible to recommend any specific option. Those who find such broad statements disappointing should remember that the panel can only judge on the basis of clinical experience and published evidence. Often, a treatment is widely held to be evidence based, although not a single study has ever been performed.

Therefore, one of our aims was to define some implications for future research. A fair amount of RCTs were retrievable to answer the various issues. We consider it unlikely that important studies were missed by our literature searches because we combined various techniques to capture all relevant studies. However, the available studies mostly focused on those questions which can be answered already using a small sample size and short-term observation. Some other statements did not receive grade A because the existing RCTs had methodological flaws. What is of general concern is that such a large proportion of trials assessed pathophysio-

logical rather than clinical outcomes. These trials, albeit randomized, are usually insufficient to answer the clinical questions we had posed. Therefore, the plea for clinically relevant RCTs cannot be reiterated too often. It is a future task to check whether the recommendations have to be modified on the basis of new data.

Developing guidelines is only worthwhile if they are used clinically. Guideline use hinges upon guideline awareness and knowledge. Therefore, the format and dissemination of the current guideline goes beyond simple publication. Since guidelines created on a European level cannot address the local circumstances in every European country or even hospital, the E.A.E.S scientific committee recommends the use of the current guideline as a basis for a locally adapted and translated guideline, which could then be implemented at any given level.

The most important factors that have to be considered before adapting this guideline for local use are individual surgical expertise and health care setting. Some surgical techniques that are discussed or even recommended here are probably not practical or affordable for every European surgeon. This is why we decided not to include cost comparisons in this guideline.

References

1. Abe K, Hashimoto N, Taniguchi A, Yoshiya I (1998) Middle cerebral artery blood flow velocity during laparoscopic surgery in head-down position. *Surg Laparosc Endosc* 8: 1–4
2. Aitola P, Airo I, Kaukinen S, Ylitalo P (1998) Comparison of N₂O and CO₂ pneumoperitoneums during laparoscopic cholecystectomy with special reference to postoperative pain. *Surg Laparosc Endosc* 8: 140–144
3. Alexander DJ, Ngoi SS, Lee L, So J, Mak K, Chan S, Goh PM (1996) Randomized trial of periportal peritoneal bupivacaine for pain relief after laparoscopic cholecystectomy. *Br J Surg* 83: 1223–1225
4. Alexander JI, Hull MGR (1987) Abdominal pain after laparoscopy: the value of a gas drain. *Br J Obstet Gynaecol* 94: 267–269
5. Alijani A, Hanna GB, Cuschieri A (2001) Cardiac function during conventional pneumoperitoneum versus mechanical abdominal wall lift in laparoscopic cholecystectomy [abstract] *Br J Surg* 88: 743–744
6. Alishahi S, Francis N, Crofts S, Duncan L, Bickel A, Cuschieri A (2001) Central and peripheral adverse hemodynamic changes during laparoscopic surgery and their reversal with a novel intermittent sequential pneumatic compression device. *Ann Surg* 233: 176–182
7. Altarac S, Janetschek G, Eder E, Bartsch G (1996) Pneumothorax complicating laparoscopic ureterolysis. *J Laparoendosc Surg* 6: 193–196
8. Amos JD, Schorr SJ, Norman PF, Poole GV, Thomae KR, Mancino AT, Hall TJ, Scott-Conner CEH (1996) Laparoscopy during pregnancy. *Am J Surg* 171: 435–437
9. Andreoli M, Servakov M, Meyers P, Mann Jr. WJ (1999) Laparoscopic surgery during pregnancy. *J Am Assoc Gynecol Laparosc* 6: 229–233
10. Baca I, Schneider B, Köhler T, Misselwitz F, Zehle A, Mühe F (1997) Thromboembolieprophylaxe bei minimal-invasiven Eingriffen und kurzstationärer Behandlung. Ergebnisse einer, multi-centrischen, prospektiven, randomisierten, kontrollierten Studie mit einem niedermolekularen Heparin. *Chirurg* 68: 1275–1280
11. Bäcklund M, Kellokumpu I, Scheinin T, von Smitten K, Tikkanen I, Lindgren L (1998) Effect of temperature of insufflated CO₂ during and after prolonged laparoscopic surgery. *Surg Endosc* 12: 1126–1130
12. Beebe DS, McNeven MP, Crain JM, Letourneau JG, Belani KG, Abrams JA, Goodale RL (1993) Evidence of venous stasis after

- abdominal insufflation for laparoscopic cholecystectomy. *Surg Gynecol Obstet* 176: 443–447
13. Bellón JM, Manzano L, Bernardos L, Ga-Honduvilla N, Larrad A, Buján J, Alvarez-Mon M (1997) Cytokine levels after open and laparoscopic cholecystectomy. *Eur Surg Res* 29: 27–34
 14. Bemelman WA, Dunker MS, Busch OR, Den Boer KT, de Wit LT, Gouma DJ (2000) Efficacy of establishment of pneumoperitoneum with the Verres needle, Hasson trocar, and modified blunt trocar (TrocDoc): a randomized study. *J Laparoendosc Adv Surg Tech A* 10: 325–330
 15. Ben-Haim M, Mandeli J, Friedman RL, Rosenthal RJ (2000) Mechanisms of systemic hypertension during acute elevation of intraabdominal pressure. *J Surg Res* 91: 101–105
 16. Berberoglu M, Dilek ON, Ercan F, Kati I, Özmen M (1998) The effect of CO₂ insufflation rate on the postlaparoscopic shoulder pain. *J Laparoendosc Adv Surg Tech A* 8: 273–277
 17. Berggren U, Gordh T, Grama D, Haglund U, Rastad J, Arvidsson D (1994) Laparoscopic versus open cholecystectomy: hospitalization, sick leave, analgesia and trauma responses. *Br J Surg* 81: 1362–1365
 18. Bessa X, Castells A, Lacy AM, Elizalde I, Delgado S, Boix L, Piñoi V, Pellisé M, García-Valdecasas JC, Piqué JM (2001) Laparoscopic-assisted vs open colectomy for colorectal cancer: influence on neoplastic cell mobilization. *J Gastrointest Surg* 5: 66–73
 19. Bhojryul S, Payne J, Steffes B, Swanstrom L, Way LW (2000) A randomized prospective study of radially expanding trocars in laparoscopic surgery. *J Gastrointest Surg* 4: 392–397
 20. Bisgaard T, Kehlet H, Rosenberg J (2001) Pain and convalescence after laparoscopic cholecystectomy. *Eur J Surg* 167: 84–96
 21. Bisgaard T, Klarskov B, Kristiansen VB, Callesen T, Schulze S, Kehlet H, Rosenberg J (1999) Multi-regional local anesthetic infiltration during laparoscopic cholecystectomy in patients receiving prophylactic multi-modal analgesia: a randomized, double-blind, placebo-controlled study. *Anesth Analg* 89: 1017–1024
 22. Bisgaard T, Klarskov B, Trap R, Kehlet H, Rosenberg J (2000) Pain after microlaparoscopic cholecystectomy. A randomized double-blind controlled study. *Surg Endosc* 14: 340–344
 23. Bloechle C, Emmermann A, Strate T, Scheurle UJ, Schneider C, Achilles E, Wolf M, Mack D, Zornig C, Broelsch CE (1998) Laparoscopic vs open repair of gastric perforation and abdominal lavage of associated peritonitis in pigs. *Surg Endosc* 12: 212–218
 24. Bloechle C, Emmermann A, Treu H, Achilles E, Mack D, Zornig C, Broelsch CE (1995) Effect of a pneumoperitoneum on the extent and severity of peritonitis induced by gastric ulcer perforation in the rat. *Surg Endosc* 9: 898–901
 25. Bloechle C, Kluth D, Holstein AF, Emmermann A, Strate T, Zomig C, Izbicki JR (1999) A pneumoperitoneum perpetuates severe damage to the ultrastructural integrity of parietal peritoneum in gastric perforation-induced peritonitis in rats. *Surg Endosc* 13: 683–688
 26. Böhm B, Junghans T, Neudecker J, Schwenk W (1999) Leber und Nierenfunktion nach laparoskopischer und konventioneller Resektion kolorektaler Tumore – Ergebnisse aus einer prospektiv randomisierten Studie. *Visceralchirurgie* 34: 20–24
 27. Böhm B, Schwenk W, Junghans T (2000) *Das pneumoperitoneum* Springer-Verlag, Berlin, pp 42–46
 28. Bongard FS, Pianim NA, Leighton TA, Dubecz S, Davis IP, Lippmann M, Klein S, Liu SY (1993) Helium insufflation for laparoscopic operation. *Surg Gynecol Obstet* 177: 140–146
 29. Bonjer HJ, Hazebroek EJ, Kazemier G, Giuffrida MC, Meijer WS, Lange JF (1997) Open versus closed establishment of pneumoperitoneum in laparoscopic surgery. *Br J Surg* 84: 599–602
 30. Borgatta L, Gruss L, Barad D, Kaali SG (1990) Direct trocar insertion vs Verres needle use for laparoscopic sterilization. *J Reprod Med* 35: 891–894
 31. Bounameaux H, Didier D, Polat O, Desmarais S, de Moerloose P, Huber O (1997) Antithrombotic prophylaxis in patients undergoing laparoscopic cholecystectomy. *Thromb Res* 86: 271–273
 32. Bouvy ND, Giuffrida MC, Tseng LN, Steyerberg EW, Marquet RL, Jeekel H, Bonjer HJ (1998) Effects of carbon dioxide pneumoperitoneum, air pneumoperitoneum, and gasless laparoscopy on body weight and tumor growth. *Arch Surg* 133: 652–656
 33. Bouvy ND, Marquet RL, Jeekel H, Bonjer HJ (1996) Impact of gas(less) laparoscopy and laparotomy on peritoneal tumor growth and abdominal wall metastases. *Ann Surg* 224: 694–701
 34. Braumann C, Jacobi CA, Ordemann J, Stöblein R, Müller JW (2000) Einfluß der intraoperativen, intravenösen und intraperitonealen Gabe von Taurolidin/Heparin in der laparoskopischen Chirurgie auf das intra- und - xtraperitoneale Tumorwachstum. In: Encke A (ed) *Chirurgisches Forum 2000 für experimentelle und klinische Forschung*, Band, pp 29,
 35. Bresadola F, Pasqualucci A, Donini A, Chiarandini P, Anania G, Terroso G, Sistu MA, Pasetto A (1999) Elective transumbilical compared with standard laparoscopic cholecystectomy. *Eur J Surg* 165: 29–34
 36. Bruce DM, Smith M, Walker CB, Heys SD, Binnie NR, Gough DB, Broom J, Eremin O (1999) Minimal access surgery for cholelithiasis induces an attenuated acute phase response. *Am J Surg* 178: 232–234
 37. Buchmann P, Christen D, Moll C, Flury R, Sartoretti C (1996) Tumorzellen in der peritonealen Spülflüssigkeit bei konventionellen und laparoskopischen Operationen wegen kolorektaler Karzinome. *Swiss Surg Suppl* 4: 45–49
 38. Byrne J, Timon D, Armstrong C, Horgan PG, Quill DS (1994) A comparison of analgesic requirements and pulmonary function in open versus laparoscopic cholecystectomy. *Minimally Invasive Ther* 3: 3–6
 39. Byron JW, Markenson G, Miyazawa K (1993) A randomized comparison of Verres needle and direct trocar insertion for laparoscopy. *Surg Gynecol Obstet* 177: 259–262
 40. Callesen T, Hjort D, Mogensen T, Schouenborg L, Nielsen D, Reventlid H, Kehlet H (1999) Combined field block and i.p. instillation of ropivacaine for pain management after laparoscopic sterilization. *Br J Anaesth* 82: 586–590
 41. Caprini JA, Arcelus JI, Laubach M, Size G, Hoffman KN, Coats RW, Blattner S (1995) Postoperative hypercoagulability and deep-vein thrombosis after laparoscopic cholecystectomy. *Surg Endosc* 9: 304–309
 42. Casati A, Valentini G, Ferrari S, Senatore R, Zangrillo A, Torri G (1997) Cardiorespiratory changes during gynaecological laparoscopy by abdominal wall elevation: comparison with carbon dioxide pneumoperitoneum. *Br J Anaesth* 78: 51–54
 43. Cason CL, Seidel SL, Bushmaier M (1996) Recovery from laparoscopic cholecystectomy procedures. *AORN J* 63: 1099–1116
 44. Chandler JG, Corson SL, Way LW (2001) Three spectra of laparoscopic entry access injuries. *J Am Coll Surg* 192: 478–491
 45. Chaudhary D, Verma GR, Gupta R, Bose SM, Ganguly NK (1999) Comparative evaluation of the inflammatory mediators in patients undergoing laparoscopic versus conventional cholecystectomy. *Aust N Z J Surg* 69: 369–372
 46. Cheah WK, Lenzi JE, So JBY, Kum CK, Goh PMY (2001) Randomized trial of needleless versus laparoscopic cholecystectomy. *Br J Surg* 88: 45–47
 47. Chekan EG, Nataraj C, Clary EM, Haywati TZ, Brody FJ, Stamat JC, Fina MC, Eubanks WS, Westcott CJ (1999) Intraperitoneal immunity and pneumoperitoneum. *Surg Endosc* 13: 1135–1138
 48. Chumillas S, Ponce JL, Delgado F, Viciano V (1998) Pulmonary function and complications after laparoscopic cholecystectomy. *Eur J Surg* 164: 433–437
 49. Chundrigar T, Hedges AR, Morris R, Stamatakis JD (1993) Intraperitoneal bupivacaine for effective pain relief after laparoscopic cholecystectomy. *Ann R Coll Surg England* 75: 437–439
 50. Cogliandolo A, Manganaro T, Saitta FP, Micali B (1998) Blind versus open approach to laparoscopic cholecystectomy: a randomized study. *Surg Laparosc Endosc* 8: 353–355
 51. Coskun I, Hatipoglu AR, Topaloglu A, Yoruk Y, Yalcinkaya S, Caglar T (2000) Laparoscopic versus open cholecystectomy: effect on pulmonary function tests. *Hepatogastroenterology* 47: 341–342
 52. Cottin V, Delafosse B, Viale JP (1996) Gas embolism during laparoscopy: a report of seven cases in patients with previous abdominal surgical history. *Surg Endosc* 10: 166–169
 53. Cravello L, D'Ercole C, Roger V, Samson D, Blanc B (1999) Laparoscopic surgery in gynecology: randomized prospective study comparing pneumoperitoneum and abdominal wall suspension. *Eur J Obstet Gynecol Reprod Biol* 83: 9–14
 54. Cruz AM, Southerland LC, Duke T, Townsend HG, Ferguson JG, Crone LA (1996) Intraabdominal carbon dioxide insufflation in the pregnant ewe. Uterine blood flow, intraamniotic pressure, and cardiopulmonary effects. *Anesthesiology* 85: 1395–1402

55. Cunniffe MG, McAnena OJ, Dar MA, Calleary J, Flynn N (1998) A prospective randomized trial of intraoperative bupivacaine irrigation for management of shoulder-tip pain following laparoscopy. *Am J Surg* 176: 258–261
56. Curet MJ, Allen D, Josloff RK, Pitcher DE, Curet LB, Miscall BG, Zucker KA (1996) Laparoscopy during pregnancy. *Arch Surg* 131: 546–551
57. Curet MJ, Putrakul K, Pitcher DE, Josloff RK, Zucker KA (2000) Laparoscopically assisted colon resection for colon carcinoma: perioperative results and long-term outcome. *Surg Endosc* 14: 1062–1066
58. Curet MJ, Vogt DA, Schob O, Quails C, Izquierdo LA, Zucker KA (1996) Effects of CO₂ pneumoperitoneum in pregnant ewes. *J Surg Res* 63: 339–344
59. Curet MJ, Weber DM, Sae A, Lopez J (2001) Effects of helium pneumoperitoneum in pregnant ewes. *Surg Endosc* 15: 710–714
60. De Cosmo G, Iannace E, Primieri P, Valente MR, Proietti R, Matteis M, Silvestrini M (2001) Changes in cerebral hemodynamics during laparoscopic cholecystectomy. *Neurol Res* 21: 658–660
61. Derouin M, Couture P, Boudreault D, Girard D, Gravel D (1996) Detection of gas embolism by transesophageal echocardiography during laparoscopic cholecystectomy. *Anesth Analg* 82: 119–124
62. de Wilde RL (1991) Goodbye to late bowel obstruction after appendectomy. *Lancet* 338: 1012
63. Dexter SP, Vucevic M, Gibson J, McMahon MJ (1999) Hemodynamic consequences of high- and low-pressure capnoperitoneum during laparoscopic cholecystectomy. *Surg Endosc* 13: 376–381
64. Dhoste K, Lacoste L, Karayan J, Lehuede MS, Thomas D, Fusciardi J (1996) Haemodynamic and ventilatory changes during laparoscopic cholecystectomy in elderly ASA III patients. *Can J Anaesth* 43: 783–788
65. Dion YM, Levesque C, Doillon CJ (1995) Experimental carbon dioxide pulmonary embolization after vena cava laceration under pneumoperitoneum. *Surg Endosc* 9: 1065–1069
66. Di Vita G, Sciume C, Lauria G, Stella C, Raimondo D, Leo P (2000) II sistema fibrinolitico dopo colecistectomia laparoscopica. *Minerva Chir* 55: 587–592
67. Drummond GB, Martin LV (1978) Pressure–volume relationships in the lung during laparoscopy. *Br J Anaesth* 50: 261–270
68. Eleftheriadis E, Kotzampassi K, Botsios D, Tzartinglou E, Farmakis H, Dadoukis J (1996) Splanchnic ischemia during laparoscopic cholecystectomy. *Surg Endosc* 10: 324–326
69. Elhakim M, Amine H, Kamel S, Saad F (2000) Effects of intraperitoneal lidocaine combined with intravenous or intraperitoneal toxicam on pain relief and bowel recovery after laparoscopic cholecystectomy. *Acta Anaesthesiol Scand* 44: 929–933
70. Elhakim M, Elkott M, Ali NM, Tahoun HM (2000) Intraperitoneal lidocaine for postoperative pain after laparoscopy. *Acta Anaesthesiol Scand* 44: 280–284
71. Elliott DC, Rodriguez A, Moncure M, Myers RA, Shillinglaw W, Davis F, Goldberg A, Mitchell K, McRitchie D (1998) The accuracy of diagnostic laparoscopy in trauma patients: a prospective, controlled study. *Int Surg* 83: 294–298
72. Ellström M, Bengtsson A, Tylman M, Haeger M, Olsson JH, Hahlin M (1996) Evaluation of tissue trauma after laparoscopic and abdominal hysterectomy: measurements of neutrophil activation and release of interleukin-6, cortisol, and C-reactive protein. *J Am Coll Surg* 182: 423–430
73. Ellström M, Olsén MF, Olsson JH, Nordberg G, Bengtsson A, Hahlin M (1998) Pain and pulmonary function following laparoscopic and abdominal hysterectomy: a randomized study. *Acta Obstet Gynecol Scand* 77: 923–928
74. Emeljanov SI, Fedenko W, Levite EM, Panfilov SA, Bobrinskaya IG, Fedorov AV, Matveev NL, Evdoshenko W, Luosev SV, Bokarev W, Musaeva SR (1998) Pneumoperitoneum risk prognosis and correction of venous circulation disturbances in laparoscopic surgery. A pilot study. *Surg Endosc* 12: 1224–1231
75. Este-McDonald JR, Josephs LG, Birkett DH, Hirsch EF (1995) Changes in intracranial pressure associated with apneic retractors. *Arch Surg* 130: 362–366
76. Evasovich MR, Clark TC, Horattas MC, Holda S, Treen L (1996) Does pneumoperitoneum during laparoscopy increase bacterial translocation? *Surg Endosc* 10: 1176–1179
77. Fabian TC, Croce MA, Stewart RM, Pritchard FE, Minard G, Kudsk KA (1993) A prospective analysis of diagnostic laparoscopy in trauma. *Ann Surg* 217: 557–565
78. Fahy BG, Hasnain JU, Flowers JL, Plotkin JS, Odonkor P, Ferguson MK (1999) Transesophageal echocardiographic detection of gas embolism and cardiac valvular dysfunction during laparoscopic nephrectomy. *Anesth Analg* 88: 500–504
79. Feig BW, Berger DH, Dougherty TB, Dupuis JF, Hsi B, Hickey RC, Ota DM (1994) Pharmacologic intervention can reestablish baseline hemodynamic parameters during laparoscopy. *Surgery* 116: 733–741
80. Feste JR, Bojahr B, Turner DJ (2000) Randomized trial comparing a radially expandable needle system with cutting trocars. *JLS* 4: 11–45
81. Fleisher LA, Yee K, Lillemo KD, Talamini MA, Yeo CJ, Heath R, Bass E, Snyder DS, Parker SD (1999) Is outpatient laparoscopic cholecystectomy safe and cost-effective? A model to study transition of care. *Anesthesiology* 90: 1746–1755
82. Fleming RY, Dougherty TB, Feig BW (1997) The safety of helium for abdominal insufflation. *Surg Endosc* 11: 230–234
83. Frank SM, Fleisher LA, Breslow MJ, Higgins MS, Olson KF, Kelly S, Beattie C (1997) Perioperative maintenance of normothermia reduces the incidence of morbid cardiac events. A randomized clinical trial. *J Am Med Assoc* 277: 1127–1134
84. Frank SM, Higgins MS, Breslow MJ, Fleisher LA, Gorman RB, Sitzmann JV, Raff H, Beattie C (1995) The catecholamine, cortisol, and hemodynamic responses to mild perioperative hypothermia. A randomized clinical trial. *Anesthesiology* 82: 83–93
85. Frank SM, Higgins MS, Fleisher LA, Sitzmann JV, Raff H, Breslow MJ (1997) Adrenergic, respiratory, and cardiovascular effects of core cooling in humans. *Am J Physiol* 272: R557–R562
86. Fredman B, Jedeikin R, Olsfanger D, Flor P, Gruzman A (1994) Residual pneumoperitoneum: a cause of postoperative pain after laparoscopic cholecystectomy. *Anesth Analg* 79: 152–154
87. Fuhrer Y, Charpentier C, Boulanger G, Menu N, Grosdidier G, Laxenaire MC (1996) Analgésie après cholécystectomie par voie coelioscopique par administration intrapéritonéale de bupivacaine. *Ann Fr Anesth Reanim* 15: 128–134
88. Fujii Y, Tanaka H, Tsuruoka S, Toyooka H, Amaha K (1994) Middle cerebral arterial blood flow velocity increases during laparoscopic cholecystectomy. *Anesth Analg* 78: 80–83
89. Fukushima R, Kawamura YJ, Saito H, Saito Y, Hashiguchi Y, Sawada T, Muto T (1996) Interleukin-6 and stress hormone responses after uncomplicated gasless laparoscopic-assisted and open sigmoid colectomy. *Dis Colon Rectum* 39: S29–S34
90. Galizia G, Prizio G, Lieto E, Castellano P, Pelosio L, Imperatore V, Ferrara A, Pignatelli C (2001) Hemodynamic and pulmonary changes during open, carbon dioxide pneumoperitoneum and abdominal wall-lifting cholecystectomy. *Surg Endosc* 15: 477–483
91. Gándara MV, de Vega DS, Escríu N, Olmedilla C, Pérez-Mencia MT, Zueras R, López A (1997) Alteraciones respiratorias durante la colecistectomía laparoscópica. Estudio comparativo de tres técnicas anestésicas. *Rev Esp Anestesiología Reanim* 44: 177–181
92. Gehring H, Kuhmann K, Klotz KF, Ocklitz E, Roth-Isigkeit A, Sedemund-Adib B, Schmucker P (1998) Effects of propofol vs isoflurane on respiratory gas exchange during laparoscopic cholecystectomy. *Acta Anaesthesiol Scand* 42: 189–194
93. Gharaibeh KL, Al-Jaberi TM (2000) Bupivacaine instillation into gallbladder bed after laparoscopic cholecystectomy: does it decrease shoulder pain? *J Laparoendosc Adv Surg Tech A* 10: 137–141
94. Giraudo G, Brachet Contul R, Caccetta M, Morino M, Glasgou RE, Visser BC, Harris HW, Patti MG, Kilpatrick SJ, Mulvihill SJ (2001) Gasless laparoscopy could avoid alterations in hepatic function. Changing management of gallstone disease during pregnancy. *Surg Endosc* 15: 741241–746246
95. Glasgou RE, Visser BC, Harris HW, Patti MG, Kilpatrick SJ, Mulvihill SJ (1998) Changing management of gallstone disease during pregnancy. *Surg Endosc* 12: 241–246
96. Goldberg JM, Maurer WG (1997) A randomized comparison of gasless laparoscopy and CO₂ pneumoperitoneum. *Obstet Gynecol* 90: 416–420
97. Golder M, Rhodes M (1998) Prospective randomized trial of 5- and 10-mm epigastric ports in laparoscopic cholecystectomy. *Br J Surg* 85: 1066–1067

98. Goodale RL, Beebe DS, McNevin MP, Boyle M, Letourneau JG, Abrams JH, Cerra FB (1993) Hemodynamic, respiratory, and metabolic effects of laparoscopic cholecystectomy. *Am J Surg* 166: 533–537
99. Gouldman JW, Sticca RP, Rippon MB, McAlhany Jr JC (1998) Laparoscopic cholecystectomy in pregnancy. *Am Surg* 64: 93–98
100. Greif WM, Forse RA (1998) Hemodynamic effects of the laparoscopic pneumoperitoneum during sepsis in a porcine endotoxemic shock model. *Ann Surg* 227: 474–480
101. Gueugniaud PY, Bertin-Maghit M, Petit P, Muchada R (1995) Diagnostic par surveillance du débit aortique et du capnogramme d'un arrêt circulatoire par embolie de CO₂ au cours de la chirurgie laparoscopique. *Ann Fr Anesth Reanim* 14: 417–420
102. Guido RS, Brooks K, McKenzie R, Gruss J, Krohn MA (1998) A randomized, prospective comparison of pain after gasless laparoscopy and traditional laparoscopy. *J Am Assoc Gynecol Laparosc* 5: 148–153
103. Gulla N, Patriti A, Lazzarini F, Tristaino B (2000) Accesso anteriore transperitoneale nella surrenalectomia videolaparoscopica. *Minerva Chir* 55: 371–375
104. Gurtner GC, Robertson CS, Chung SC, Ling TK, Ip SM, Li AK (1995) Effect of carbon dioxide pneumoperitoneum on bacteremia and endotoxaemia in an animal model of peritonitis. *Br J Surg* 82: 844–848
105. Halverson A, Buchanan R, Jacobs L, Shayani V, Hunt T, Riedel C, Sackier J (1998) Evaluation of mechanism of increased intracranial pressure with insufflation. *Surg Endosc* 12: 266–269
106. Halverson AL, Barrett WL, Iglesias AR, Lee WT, Garber SM, Sackier JM (1999) Decreased cerebrospinal fluid absorption during abdominal insufflation. *Surg Endosc* 13: 797–800
107. Harris SN, Ballantyne GH, Luther MA, Perrino AC (1996) Alterations of cardiovascular performance during laparoscopic colectomy: a combined hemodynamic and echocardiographic analysis. *Anesth Analg* 83: 482–487
108. Hashizume M, Sugimachi K (1997) Needle and trocar injury during laparoscopic surgery in Japan. *Surg Endosc* 11: 1198–1201
109. Hasson HM (1971) A modified instrument and method for laparoscopy. *Am J Obstet Gynecol* 110: 886–870
110. Hein HAT, Joshi GP, Ramsay MAE, Fox LG, Gaway BJ, Hellman CL, Arnold JC (1997) Hemodynamic changes during laparoscopic cholecystectomy in patients with severe cardiac disease. *J Clin Anesth* 9: 261–265
111. Hendolin HI, Pääkkönen ME, Alhava EM, Tarvainen R, Kemppinen T, Lahtinen P (2000) Laparoscopic or open cholecystectomy: a prospective randomised trial to compare postoperative pain, pulmonary function, and stress response. *Eur J Surg* 166: 394–399
112. Hewitt PM, Ip SM, Kwok SPY, Somers SS, Li K, Leung KL, Lau WY, Li AKC (1998) Laparoscopic-assisted vs open surgery for colorectal cancer: comparative study of immune effects. *Dis Colon Rectum* 41: 901–909
113. Hirvonen EA, Nuutinen LS, Kauko M (1995) Hemodynamic changes due to Trendelenburg positioning and pneumoperitoneum during laparoscopic hysterectomy. *Acta Anaesthesiol Scand* 39: 949–955
114. Hirvonen EA, Nuutinen LS, Vuolteenaho O (1997) Hormonal responses and cardiac filling pressures in head-up or head-down position and pneumoperitoneum in patients undergoing operative laparoscopy. *Br J Anaesth* 78: 128–133
115. Hirvonen EA, Poikolainen EO, Pääkkönen ME, Nuutinen LS (2000) The adverse hemodynamic effects of anesthesia, head-up tilt, and carbon dioxide pneumoperitoneum during laparoscopic cholecystectomy. *Surg Endosc* 14: 272–277
116. Holmdahl L, Eriksson E, Eriksson BI, Risberg B (1998) Depression of peritoneal fibrinolysis during operation is a local response to trauma. *Surgery* 123: 539–544
117. Holthausen UH, Razek TSA, Hinchey EJ, Oung CM, Chiu RC-J, Nagelschmidt M, Troidl H (1997) Monitoring des intrakraniellen Druckes im Modell eines "Schädeltraumas": Kohlendioxid (CO₂)-Pneumoperitoneum versus Laparolift-Verfahren. *Langenbecks Arch Chir Kongressbd* 114: 257–260
118. Horzic M, Korusic A, Bunoza D, Maric K (1998) The influence of increased intra-abdominal pressure on blood coagulation values. *Hepatogastroenterology* 45: 1519–1521
119. Howells GA, Uzieblo MR, Bair H, Boyer MD (2000) Tension pneumopericardium during laparoscopy for trauma. *Surg Laparosc Endosc Percutan Tech* 10: 44–46
120. Hunter JG, Swanstrom L, Thornburg K (1995) Carbon dioxide pneumoperitoneum induces fetal acidosis in a pregnant ewe model. *Surg Endosc* 9: 272–279
121. Hynes SR, Marshall RL (1992) Venous gas embolism during gynaecological laparoscopy. *Can J Anaesth* 39: 748–749
122. Ido K, Suzuki T, Kimura K, Taniguchi Y, Kawamoto C, Isoda N, Nagamine N, Ioka T, Kumagai M, Hirayama Y (1995) Lower extremity venous stasis during laparoscopic cholecystectomy as assessed using color Doppler ultrasound. *Surg Endosc* 9: 310–313
123. Irgau I, Koyfman Y, Tikellis JI (1995) Elective intraoperative intracranial pressure monitoring during laparoscopic cholecystectomy. *Arch Surg* 130: 1011–1013
124. Ishizaki Y, Bandai Y, Shimomura K, Abe H, Ohtomo Y, Idezuki Y (1993) Safe intraabdominal pressure of carbon dioxide pneumoperitoneum during laparoscopic surgery. *Surgery* 114: 549–554
125. Ishizuka B, Kuribayashi Y, Kobayashi Y, Hamada N, Abe Y, Amemiya A, Aoki T, Satoh T (2000) Stress response during laparoscopy with CO₂ insufflation and with mechanical elevation of the abdominal wall. *J Am Assoc Gynecol Laparosc* 3: 363–371
126. Iwase K, Takenaka H, Yagura A, Ishizaka T, Ohata T, Takagaki M, Oshima S (1992) Hemodynamic changes during laparoscopic cholecystectomy in patients with heart disease. *Endoscopy* 24: 771–773
127. Jackman SV, Weingart JD, Kinsman SL, Docimo SG (2000) Laparoscopic surgery in patients with ventriculoperitoneal shunts: safety and monitoring. *J Urol* 164: 1352–1354
128. Jackson SA, Laurence AS, Hill JC (1996) Does post-laparoscopy pain relate to residual carbon dioxide? *Anaesthesia* 51: 485–487
129. Jacobi CA, Ordemann J, Böhm B, Zieren HU, Volk HD, Lorenz W, Halle E, Müller JM (1997) Does laparoscopy increase bacteremia and endotoxemia in a peritonitis model? *Surg Endosc* 11: 235–238
130. Jacobi CA, Ordemann J, Zieren HU, Volk HD, Bauhofer A, Halle E, Müller JM (1998) Increased systemic inflammation after laparotomy vs laparoscopy in an animal model of peritonitis. *Arch Surg* 133: 258–262
131. Jacobi CA, Sabat R, Ordemann J, Wenger F, Volk HD, Müller JM (1997) Peritoneale Instillation von Taurolidin und Heparin zur Verhinderung von intraperitonealem Tumorstadium und Trocarkarmetastasen bei laparoskopischen Operationen im Rattenmodell. *Langenbecks Arch Chir* 382: S31–S36
132. Jacobi CA, Sterzel A, Braumann C, Halle E, Stösslein R, Krähnenbühl L, Müller JM (2001) The impact of conventional and laparoscopic colon resection (CO₂ or helium) on intraperitoneal adhesion formation in a rat peritonitis model. *Surg Endosc* 15: 380–386
133. Jacobi CA, Wenger F, Sabat R, Volk T, Ordemann J, Müller JM (1998) The impact of laparoscopy with carbon dioxide versus helium on immunologic function and tumor growth in a rat model. *Dig Surg* 15: 110–116
134. Jakeways MSR, Mitchell V, Hashim IA, Chadwick SJD, Shenkin A, Green CJ, Carri F (1994) Metabolic and inflammatory responses after open or laparoscopic cholecystectomy. *Br J Surg* 81: 127–131
135. Jakimowicz J, Stultiens G, Smulders F (1998) Laparoscopic insufflation of the abdomen reduces portal venous flow. *Surg Endosc* 12: 129–132
136. Johnson PL, Sibert KS (1997) Laparoscopy. Gasless vs CO₂ pneumoperitoneum. *J Reprod Med* 42: 255–259
137. Jorgensen JO, Gillies RB, Hunt DR, Caplehom JRM, Lumley T (1995) A simple and effective way to reduce postoperative pain after laparoscopic cholecystectomy. *Aust N Z J Surg* 65: 466–469
138. Jorgensen JO, Lalak NJ, North L, Hanel K, Hunt DR, Morris DL (1994) Venous stasis during laparoscopic cholecystectomy. *Surg Laparosc Endosc* 4: 128–133
139. Joris J, Cigarini I, Legrand M, Jacquet N, De Groote D, Franchimont P, Lamy M (1992) Metabolic and respiratory changes after cholecystectomy performed via laparotomy or laparoscopy. *Br J Anaesth* 69: 341–345
140. Joris J, Thiry E, Paris P, Weerts J, Lamy M (1995) Pain after laparoscopic cholecystectomy: characteristics and effect of intraperitoneal bupivacaine. *Anesth Analg* 81: 379–384

141. Joris JL, Chiche JD, Canivet JLM, Jacquet NJ, Legros JJ, Lamy ML (1998) Hemodynamic changes induced by laparoscopy and their endocrine correlates: effects of clonidine. *J Am Coll Cardiol* 32: 1389–1396
142. Josephs LG, Este-McDonald JR, Birkett DH, Hirsch EF (1994) Diagnostic laparoscopy increases intracranial pressure. *J Trauma* 36: 815–819
143. Joshi GP (2001) Complications of laparoscopy. *Anesthesiol Clin North Am* 19: 89–105
144. Junghans T, Böhm B, Gründel K, Scheiba-Zórron R, Müller JM (1999) Effects of induced intravenous helium and CO₂ embolism on the cardiovascular system. *Minimal Invasive Chirurgie* 8: 52–56
145. Junghans T, Böhm B, Gründel K, Schwenk W, Müller JM (1997) Does pneumoperitoneum with different gases, body positions, and intraperitoneal pressures influence renal and hepatic blood flow? *Surgery* 121: 206–211
146. Junghans T, Böhm B, Meyer E (2000) Influence of nitrous oxide anesthesia on venous embolism with carbon dioxide and helium during pneumoperitoneum. *Surg Endosc* 14: 1167–1170
147. Junghans T, Böhm B, Neudecker J, Mansmann U, Gründel K (1999) Auswirkungen von Argon-Gasembolien während eines Pneumoperitoneums. *Chirurg* 70: 184–189
148. Kanellos I, Zaroglidis K, Ziovas E, Dadoukis I (1994) Prospektiv-vergleichende Studie der Lungenfunktion nach laparoskopischer, Mini-Lap- oder konventioneller Cholezystectomie. *Minimal Invasive Chirurgie* 4: 169–171
149. Karayiannakis AJ, Makri GG, Mantzioka A, Karousos D, Karatzas G (1997) Systemic stress response after laparoscopic or open cholecystectomy: a randomized trial. *Br J Surg* 84: 467–471
150. Kazemier G, Berends FJ, Bouvy ND, Lange JF, Bonjer HJ (1997) The influence of pneumoperitoneum on the peritoneal implantation of free intraperitoneal cancer cells. *Surg Endosc* 11: 698–699
151. Khan AU, Pandya K, Clifton MA (1995) Near fatal gas embolism during laparoscopic cholecystectomy. *Ann R Coll Surg Engl* 77: 67–68
152. Kiviluoto T, Sirén J, Luukkonen P, Kivilaakso E (1998) Randomized trial of laparoscopic versus open cholecystectomy for acute and gangrenous cholecystitis. *Lancet* 351: 321–325
153. Knolmayer TJ, Bowyer MW, Egan JC, Asbun HJ (1998) The effects of pneumoperitoneum on gastric blood flow and traditional hemodynamic measurements. *Surg Endosc* 12: 115–118
154. Koivusalo AM, Kellokumpu I, Lindgren L (1996) Gasless laparoscopic cholecystectomy: comparison of postoperative recovery with conventional technique. *Br J Anaesth* 77: 576–580
155. Koivusalo AM, Kellokumpu I, Lindgren L (1997) Postoperative drowsiness and emetic sequelae correlate to total amount of carbon dioxide used during laparoscopic cholecystectomy. *Surg Endosc* 11: 42–44
156. Koivusalo AM, Kellokumpu I, Ristkari S, Lindgren L (1997) Splanchnic and renal deterioration during and after laparoscopic cholecystectomy: a comparison of the carbon dioxide pneumoperitoneum and the abdominal wall lift method. *Anesth Analg* 85: 886–891
157. Koivusalo AM, Kellokumpu I, Scheinin M, Tikkanen I, Halme L, Lindgren L (1996) Randomized comparison of the neuroendocrine response to laparoscopic cholecystectomy using either conventional or abdominal wall lift techniques. *Br J Surg* 83: 1532–1536
158. Koivusalo AM, Kellokumpu I, Scheinin M, Tikkanen I, Mäkisalo H, Lindgren L (1998) A comparison of gasless mechanical and conventional carbon dioxide pneumoperitoneum methods for laparoscopic cholecystectomy. *Anesth Analg* 86: 153–158
159. Koivusalo AM, Lindgren L (1999) Respiratory mechanics during laparoscopic cholecystectomy. *Anesth Analg* 89: 800
160. Koivusalo AM, Lindgren L (2000) Effects of carbon dioxide pneumoperitoneum for laparoscopic cholecystectomy. *Acta Anaesthesiol Scand* 44: 834–841
161. Koivusalo AM, Scheinin M, Tikkanen I, Yli-Suomu T, Ristkari S, Laakso J, Lindgren L (1998) Effects of esmolol on haemodynamic response to CO₂ pneumoperitoneum for laparoscopic surgery. *Acta Anaesthesiol Scand* 42: 510–517
162. Korell M, Schmaus F, Strowitzki T, Schneeweiss SG, Hepp H (1996) Pain intensity following laparoscopy. *Surg Laparosc Endosc* 6: 375–379
163. Kraut EJ, Anderson JT, Safwat A, Barbosa R, Wolfe BM (1999) Impairment of cardiac performance by laparoscopy in patients receiving positive end-expiratory pressure. *Arch Surg* 134: 76–80
164. Kristiansson M, Saraste L, Soop M, Sundqvist KG, Thörne A (1999) Diminished interleukin-6 and C-reactive protein responses to laparoscopic versus open cholecystectomy. *Acta Anaesthesiol Scand* 43: 146–152
165. Krug F, Herald A, Wenk H, Bruch HP (1995) Narbenhernien nach laparoskopischen Eingriffen. *Chirurg* 66: 419–423
166. Kum CK, Eypasch E, Aljaziri A, Troidl H (1996) Randomized comparison of pulmonary function after the “French” and “American” techniques of laparoscopic cholecystectomy. *Br J Surg* 83: 938–941
167. Kuthe A, Tamme C, Saemann T, Schneider C, Kockering F (1999) Die laparoskopische Cholezystektomie mit Mini-Instrumentarium. *Zentralbl Chir* 124: 749–753
168. Lacy AM, Delgado S, Garcia-Valdecasas JC, Castells A, Piqué JM, Grande L, Fuster J, Tararona EM, Pera M, Visa J (1998) Port site metastases and recurrence after laparoscopic colectomy. A randomized trial. *Surg Endosc* 12: 1039–1042
169. Lajer H, Widecrantz S, Heisterberg L (1997) Hernias in trocar ports following abdominal laparoscopy. A review. *Acta Obstet Gynecol Scand* 76: 389–393
170. Lam TYD, Lee SW, So HS, Kwok SPY (2000) Radially expanding trocar: a less painful alternative for laparoscopic surgery. *J Laparoendosc Adv Surg Tech A* 10: 269–273
171. Lantz PE, Smith JD (1994) Fatal carbon dioxide embolism complicating attempted laparoscopic cholecystectomy—case report and literature review. *J Forensic Sci* 39: 1468–1480
172. Lau JYW, Lo SY, Ng EKW, Lee DWH, Lam YH, Chung SCS (1998) A randomized comparison of acute phase response and endotoxemia in patients with perforated peptic ulcers receiving laparoscopic or open patch repair. *Am J Surg* 175: 325–327
173. Le Blanc-Louvry I, Coquerel A, Koning E, Maillot C, Ducrotte P (2000) Operative stress response is reduced after laparoscopic compared to open cholecystectomy: the relationship with postoperative pain and ileus. *Dig Dis Sci* 45: 1703–1713
174. Lejus C, Delile L, Plattner V, Baron M, Guillou S, Héroudy Y, Souron R (1996) Randomized, single-blinded trial of laparoscopic versus open appendectomy in children: effects on postoperative analgesia. *Anesthesiology* 84: 801–806
175. Leung KL, Lai PBS, Ho RLK, Meng WCS, Yiu RYC, Lee JFY, Lau WY (2000) Systemic cytokine response after laparoscopic-assisted resection of rectosigmoid carcinoma: A prospective randomized trial. *Ann Surg* 231: 506–511
176. Levy T, Dicker D, Shalev J, Dekel A, Farhi J, Peleg D, Ben-Rafael Z (1995) Laparoscopic unwinding of hyperstimulated ischaemic ovaries during the second trimester of pregnancy. *Hum Reprod* 10: 1478–1480
177. Lindberg F, Rasmussen I, Siegbahn A, Bergqvist D (2000) Coagulation activation after laparoscopic cholecystectomy in spite of thromboembolism prophylaxis. *Surg Endosc* 14: 858–861
178. Lindgren L, Koivusalo AM, Kellokumpu I (1995) Conventional pneumoperitoneum compared with abdominal wall lift for laparoscopic cholecystectomy. *Br J Anaesth* 75: 567–572
179. Linhares L, Jeanpierre H, Borie F, Fingerhut A, Millat B (2001) Lavage by laparoscopy fares better than lavage by laparotomy: experimental evidence. *Surg Endosc* 15: 85–89
180. Lipscomb GH, Summitt RL, McCord ML, Ling FW (1994) The effect of nitrous oxide and carbon dioxide pneumoperitoneum on operative and postoperative pain during laparoscopic sterilization under local anesthesia. *J Am Assoc Gynecol Laparosc* 2: 57–60
181. Lo CM, Liu CL, Fan ST, Lai ECS, Wong J (1998) Prospective randomized study of early versus delayed laparoscopic cholecystectomy for acute cholecystitis. *Ann Surg* 227: 461–467
182. Loeckinger A, Kleinsasser A, Hoermann C, Gassner M, Keller C, Lindner KH (2000) Inert gas exchange during pneumoperitoneum at incremental values of positive end-expiratory pressure. *Anesth Analg* 90: 466–471
183. Lord RV, Ling JJ, Hugh TB, Coleman MJ, Doust BD, Nivison-Smith I (1998) Incidence of deep vein thrombosis after laparoscopic vs minilaparotomy cholecystectomy. *Arch Surg* 133: 967–973

184. Lukban JC, Jaeger J, Hammond KC, LoBraico DA, Gordon AMC, Graebe RA (2000) Gasless versus conventional laparoscopy. *N J Med* 97: 29–34
185. Lunderoff P, Thorburn J, Hahlin M, Kallfelt B, Lindblom B (1991) Laparoscopic surgery in ectopic pregnancy. A randomized trial versus laparotomy. *Acta Obstet Gynecol Scand* 70: 343–348
186. Mäkinen MT, Heinonen PO, Klemola UM, Yli-Hankala A (2001) Gastric air tonometry during laparoscopic cholecystectomy: a comparison of two PaCO₂ levels. *Can J Anaesth* 48: 121–128
187. Mäkinen MT, Yli-Hankala A (1996) The effect of laparoscopic cholecystectomy on respiratory compliance as determined by continuous spirometry. *J Clin Anesth* 8: 119–122
188. Mall JW, Schwenk W, Rödiger O, Zippel K, Pollmann C, Müller JM (2001) Blinded prospective study of the incidence of deep venous thrombosis following conventional or laparoscopic colorectal resection. *Br J Surg* 88: 99–100
189. Marquez J, Sladen A, Gendell H, Boehnke M, Mendelow H (1981) Paradoxical cerebral air embolism without an intracardiac septal defect. Case report. *J Neurosurg* 55: 997–1000
190. Martin IG, Dexter SPL, McMahon MJ (1996) Laparoscopic cholecystectomy in pregnancy. *Surg Endosc* 10: 508–510
191. Martínez-Ramos C, Lopez-Pastor A, Nunez-Pena JR, Gopegui M, Sanz-Lopez R, Jorgensen T, Pastor L, Fernandez-Chacon JL, Tamames-Escobar S (1999) Changes in hemostasis after laparoscopic cholecystectomy. *Surg Endosc* 13: 476–479
192. Matsuda T, Ogura K, Uchida J, Fujita I, Terachi T, Yoshida O (1995) Smaller ports result in shorter convalescence after laparoscopic varicocele. *J Urol* 153: 1175–1177
193. Matsumoto T, Tsuboi S, Dolgor B, Bandoh T, Yoshida T, Kitano S (2001) The effect of gases in the intraperitoneal space on cytokine response and bacterial translocation in a rat model. *Surg Endosc* 15: 80–84
194. McDougall EM, Monk TG, Wolf JS, Hicks M, Clayman RV, Gardner S, Humphrey PA, Sharp T, Martin K (1996) The effect of prolonged pneumoperitoneum on renal function in an animal model. *J Am Coll Surg* 182: 317–328
195. McKellar DP, Anderson CT, Boynton CJ, Peoples JB (1992) Cholecystectomy during pregnancy without fetal loss. *Surg Gynecol Obstet* 174: 465–468
196. McMahon AJ, Baxter JN, Kenny G, O'Dwyer PJ (1993) Ventilatory and blood gas changes during laparoscopic and open cholecystectomy. *Br J Surg* 80: 1252–1254
197. McMahon AJ, O'Dwyer PJ, Cruikshank AM, McMillan DC, O'Reilly DS, Lowe GD, Rumley A, Logan RW, Baxter JN (1993) Comparison of metabolic responses to laparoscopic and minilaparotomy cholecystectomy. *Br J Surg* 80: 1255–1258
198. Mecke H, Kroll K (1999) Postlaparoskopisches Schmerzsyndrom nach Pneumoperitoneum mit angewärmten und kaltem CO₂-Gas. *Geburtshilfe Frauenheilkunde* 59: 611–615
199. Mehta M, Sokoll MD, Gergis SD (1984) Effects of venous air embolism on the cardiovascular system and acid base balance in the presence and absence of nitrous oxide. *Acta Anaesthesiol Scand* 28: 226–231
200. Meijer DW, Rademaker BPM, Schlooz S, Bemelman WA, de Wit LT, Bannenberg JGG, Stijnen T, Gouma DF (1997) Laparoscopic cholecystectomy using abdominal wall retraction. Hemodynamics and gas exchange, a comparison with conventional pneumoperitoneum. *Surg Endosc* 11: 645–649
201. Michaloliakou C, Chung F, Sharma S (1996) Preoperative multimodal analgesia facilitates recovery after ambulatory laparoscopic cholecystectomy. *Anesth Analg* 82: 44–51
202. Milheiro A, Sousa FC, Manso EC, Leitao F (1994) Metabolic responses to cholecystectomy: open vs laparoscopic approach. *J Laparoendosc Surg* 4: 311–317
203. Millard JA, Hill BB, Cook PS, Fenoglio ME, Stahlgren LH (1993) Intermittent sequential pneumatic compression in prevention of venous stasis associated with pneumoperitoneum during laparoscopic cholecystectomy. *Arch Surg* 128: 914–919
204. Milsom JW, Böhm B, Hammerhofer KA, Fazio V, Steiger E, Elson P (1998) A prospective, randomized trial comparing laparoscopic versus conventional techniques in colorectal cancer surgery: a preliminary report. *J Am Coll Surg* 187: 46–55
205. Mimica Z, Biocic M, Bacic A, Banovic I, Tocilj J, Radonic V, Ilic N, Petricevic A (2000) Laparoscopic and laparotomic cholecystectomy: a randomized trial comparing postoperative respiratory function. *Respiration* 67: 153–158
206. Minoli G, Terruzzi V, Spinzi GC, Benvenuti C, Rossini A (1982) The influence of carbon dioxide and nitrous oxide on pain during laparoscopy: a double-blind, controlled trial. *Gastrointest Endosc* 28: 173–175
207. Moncure M, Salem R, Moncure K, Testaiuti M, Marburger R, Ye X, Brathwaite C, Ross SE (1999) Central nervous system metabolic and physiologic effects of laparoscopy. *Am Surg* 65: 168–172
208. Morino M, Giraudo G, Festa V (1998) Alterations in hepatic function during laparoscopic surgery. An experimental clinical study. *Surg Endosc* 12: 968–972
209. Mouton WG, Bessell JR, Millard SH, Baxter PS, Maddern GJ (1999) A randomized controlled trial assessing the benefit of humidified insufflation gas during laparoscopic surgery. *Surg Endosc* 13: 106–108
210. Mouton WG, Naef M, Bessell JR, Wagner HE, Maddern GJ (2001) A randomized controlled trial to determine the effect of humidified carbon dioxide (CO₂) insufflation on postoperative pain following thoracoscopic procedures. *Surg Endosc* 15: 579–581
211. Mraovic B, Jurisic T, Kogler-Majeric V, Sustic A (1997) Intraperitoneal bupivacaine for analgesia after laparoscopic cholecystectomy. *Acta Anaesthesiol Scand* 41: 193–196
212. Myre K, Buanes T, Smith G, Stokland O (1997) Simultaneous hemodynamic and echocardiographic changes during abdominal gas insufflation. *Surg Laparosc Endosc* 7: 415–419
213. Nagelschmidt M, Holthausen U, Goost H, Fu ZX, Minor T, Troidl H, Neugebauer E (2000) Evaluation of the effects of a pneumoperitoneum with carbon dioxide or helium in a porcine model of endotoxemia. *Langenbecks Arch Surg* 385: 199–206
214. Narchi P, Benhamou D, Fernandez H (1991) Intraperitoneal local anaesthetic for shoulder pain after day-case laparoscopy. *Lancet* 338: 1569–1570
215. Nelskyla K, Yli-Hankala A, Sjöberg J, Korhonen I, Korttila K (1999) Wanning of insufflation gas during laparoscopic hysterectomy: effect on body temperature and the autonomic nervous system. *Acta Anaesthesiol Scand* 43: 974–978
216. Neudecker J, Junghans T, Ziemer H, Schwenk W (2001) Einfluß der Operationstechnik auf das fibrinolytische Potential des Peritoneums und der Peritonealflüssigkeit bei laparoskopischen und konventionellen kolorektalen Resektionen - Eine prospektiv randomisierte Studie. In: Schönleben K (ed) *Chirurgisches Forum 2001 für experimentelle und klinische Forschung*: Band 30. Springer, Berlin, pp 539
217. Neugebauer E, Sauerland S, Troidl H (2000) Recommendations for evidence-based, endoscopic surgery. The updated EAES consensus development conferences. Springer, Paris
218. Neuhaus SJ, Watson DI, Ellis T, Rofe AM, Mathew G, Jamieson GG (2000) Influence of gases on intraperitoneal immunity during laparoscopy in tumor-bearing rats. *World J Surg* 24: 1227–1231
219. Nezhat FR, Silfen SL, Evans D, Nezhat C (1991) Comparison of direct insertion of disposable and standard reusable laparoscopic trocars and previous pneumoperitoneum with Veress needle. *Obstet Gynecol* 78: 148–150
220. Ninomiya K, Kitano S, Yoshida T, Bandoh T, Baatar D, Matsumoto T (1998) Comparison of pneumoperitoneum and abdominal wall lifting as to hemodynamics and surgical stress response during laparoscopic cholecystectomy. *Surg Endosc* 12: 124–128
221. Nordentoft T, Bringstrup FA, Bremmelgaard A, Stage JG (2000) Effect of laparoscopy on bacteremia in acute appendicitis: a randomized controlled study. *Surg Laparosc Endosc Percutan Tech* 10: 302–304
222. Odeberg S, Ljungqvist O, Sollevi A (1998) Pneumoperitoneum for laparoscopic cholecystectomy is not associated with compromised splanchnic circulation. *Eur J Surg* 164: 843–848
223. Oghihara Y, Isshiki A, Kindscher JD, Goto H (1999) Abdominal wall lift versus carbon dioxide insufflation for laparoscopic resection of ovarian tumors. *J Clin Anesth* 11: 406–412
224. Ordemann J, Jacobi C, Schwenk W, Stösslein R, Müller JM (2001) Cellular and humoral inflammatory response after laparoscopic and conventional colorectal resections—results of a prospective randomized trial. *Surg Endosc* 15: 600–608

225. Ortega AE, Peters JH, Incarbone R, Estrada L, Ehsan A, Kwan Y, Spencer CJ, Moore-Jeffries E, Kuchta K, Nicoloff JT (1996) A prospective randomized comparison of the metabolic and stress hormonal responses of laparoscopic and open cholecystectomy. *J Am Coll Surg* 183: 249–256
226. Ott DE, Rejch H, Love B, McCorvey R, Toledo A, Liu CY, Syed R, Kumar K (1998) Reduction of laparoscopic-induced hypothermia, postoperative pain and recovery room length of stay by pre-conditioning gas with the Insuflow device: a prospective randomized controlled multicenter study. *JSL* 2: 321–329
227. Pasqualucci A, Contardo R, Da Broi U, Colo F, Terrosu G, Donini A, Sorrentino M, Pasetto A, Bresadola F (1994) The effects of intraperitoneal local anesthetic on analgesic requirements and endocrine response after laparoscopic cholecystectomy: a randomized double-blind controlled study. *J Laparoendosc Surg* 4: 405–412
228. Pasqualucci A, de Angelis V, Contardo R, Colo F, Terrosu G, Donini A, Pasetto A, Bresadola F (1996) Preemptive analgesia: intraperitoneal local anesthetic in laparoscopic cholecystectomy. A randomized, double-blind, placebo-controlled study. *Anesthesiology* 85: 11–20
229. Patel MI, Hardman DTA, Nicholls D, Fisher CM, Appleberg M (1996) The incidence of deep venous thrombosis after laparoscopic cholecystectomy. *Med J Aust* 164: 652–654, 656
230. Péchinot M, Rapenne T, Galloux Y, Vérain C (1996) Embolie gazeuse au cours d'une cholecystectomie par coelioscopie. *Cah Anesthesiol* 44: 365–367
231. Peitgen K, Nimitz K, Hellinger A, Walz MK (1997) Offener Zugang oder Veress-Nadel bei laparoskopischen Eingriffen? Ergebnisse einer prospektiv randomisierten Studie. *Chirurg* 68: 910–913
232. Perez JE, Alberts WM, Mamel JJ (1997) Delayed tension pneumothorax after laparoscopy. *Surg Laparosc Endosc* 7: 70–72
233. Perry CP, Tombrello R (1993) Effect of fluid instillation on postlaparoscopic pain. *J Reprod Med* 38: 768–770
234. Perttilä J, Salo M, Ovaska J, Grönroos J, Lavonius M, Katila A, Lähtenmäki M, Pulkki K (1999) Immune response after laparoscopic and conventional Nissen fundoplication. *Eur J Surg* 165: 21–28
235. Philips PA, Amaral JF (2001) Abdominal access complications in laparoscopic surgery. *J Am Coll Surg* 192: 525–536
236. Pier A, Benedic M, Mann B, Buck V (1994) Das postlaparoskopische Schmerzsyndrom: Ergebnisse einer prospektiven, randomisierten Studie. *Chirurg* 65: 200–208
237. Popesco D, Le Mière E, Maitre B, Darchy B, Domart Y (1997) Embolie gazeuse coronaire lors de la chirurgie coelioscopique. *Ann Fr Anesth Reanim* 16: 381–385
238. Popken F, Kühle R, Heintz A, Junginger T (1997) Die laparoskopische Cholecystektomie beim Hochrisikopatienten. *Chirurg* 68: 801–805
239. Popken F, Kühle R, Heintz A, Junginger T (1998) Die Cholecystektomie beim Hochrisikopatienten: Ein Vergleich zwischen konventionellem und laparoskopischen Verfahren. *Chirurg* 69: 61–65
240. Popken CA, Compton RP, Walter DN, Browder IW (1995) Benefits of pulmonary artery catheter and transesophageal echocardiographic monitoring in laparoscopic cholecystectomy patients with cardiac disease. *Am J Surg* 169: 202–207
241. Presson RG, Kirk KR, Haselby KA, Wagner WW (1991) Effect of ventilation with soluble and diffusible gases on the size of air emboli. *J Appl Physiol* 70: 1068–1074
242. Prisco D, De Gaudio AR, Caria R, Gori AM, Fedi S, Cella AP, Ginsini GF, Abbate R (2000) Videolaparoscopic cholecystectomy induces a hemostasis activation of lower grade than does open surgery. *Surg Endosc* 14: 170–174
243. Puttick MI, Scott-Coombes DM, Dye J, Nduka CC, Menzies-Gow NM, Mansfield AO, Darzi A (1999) Comparison of immunologic and physiologic effects of CO₂ pneumoperitoneum at room and body temperatures. *Surg Endosc* 13: 572–575
244. Rademaker BMP, Kalkman CJ, Odoom JA, de Wit L, Ringers J (1994) Intraperitoneal local anaesthetics after laparoscopic cholecystectomy: effects on postoperative pain, metabolic responses and lung function. *Br J Anaesth* 72: 263–266
245. Rasmussen YH, Leikersfeldt G, Drenck NE (1998) Forced-air surface warming versus oesophageal heat exchanger in the prevention of perioperative hypothermia. *Acta Anaesthesiol Scand* 42: 348–352
246. Razvi HA, Fields D, Vargas JC, Vaughan ED, Vukasin A, Sosa RE (1996) Oliguria during laparoscopic surgery: evidence for direct renal parenchymal compression as an etiologic factor. *J Endouro* 10: 1–4
247. Redmond HP, Watson RWG, Houghton T, Condon C, Watson RKG, Bouchier-Hayes D (1994) Immune function in patients undergoing open vs laparoscopic cholecystectomy. *Arch Surg* 129: 1240–1246
248. Retzke U, Graf H, Schmidt M (1998) Appendicitis in graviditate. *Zentralbl Chir* 123: 61–65
249. Rishimani ASM, Gautam SC (1996) Hemodynamic and respiratory changes during laparoscopic cholecystectomy with high and reduced intraabdominal pressure. *Surg Laparosc Endosc* 6: 201–204
250. Roberts MW, Mathiesen KA, Ho HS, Wolfe BM (1997) Cardiopulmonary responses to intravenous infusion of soluble and relatively insoluble gases. *Surg Endosc* 11: 341–346
251. Rosenthal RJ, Friedman RL, Kahn AM, Martz J, Thiagarajah S, Cohen D, Shi Q, Nussbaum M (1998) Reasons for intracranial hypertension and hemodynamic instability during acute elevations of intra-abdominal pressure: observations in a large animal model. *J Gastrointest Surg* 2: 415–425
252. Rosenthal RJ, Hiatt JR, Phillips EH, Hewitt W, Demetriou AA, Grode M (1997) Intracranial pressure. Effects of pneumoperitoneum in a large-animal model. *Surg Endosc* 11: 376–380
253. Roumen RMH, van Meurs PA, Kuypers HHC, Kraak WAG, Sauerwein RW (1992) Serum interleukin-6 and C reactive protein responses in patients after laparoscopic or conventional cholecystectomy. *Eur J Surg* 158: 541–544
254. Saad S, Minor I, Mohri T, Nagelschmidt M (2000) The clinical impact of warmed insufflation carbon dioxide gas for laparoscopic cholecystectomy. *Surg Endosc* 14: 787–790
255. Sackett DL, Straus SE, Richardson WS, Rosenberg W, Haynes RB (2000) Evidence-based medicine: How to practice and teach EBM, 2nd edn. Churchill Livingstone, London
256. Safran D, Sgambati S, Orlando R (1993) Laparoscopy in high-risk cardiac patients. *Surg Gynecol Obstet* 176: 548–554
257. Sarac AM, Aktan AO, Baykan N, Yegen C, Yalin R (1996) The effect and timing of local anesthesia in laparoscopic cholecystectomy. *Surg Laparosc Endosc* 6: 362–366
258. Sarli L, Costi R, Sansebastiano G, Trivelli M, Roncoroni L (2000) Prospective randomized trial of low-pressure pneumoperitoneum for reduction of shoulder-tip pain following laparoscopy. *Br J Surg* 87: 1161–1165
259. Sato N, Kawamoto M, Yuge O, Suyama H, Sanuki M, Matsumoto C, Inoue K (2000) Effects of pneumoperitoneum on cardiac autonomic nervous activity evaluated by heart rate variability analysis during sevoflurane, isoflurane, or propofol anesthesia. *Surg Endosc* 14: 362–366
260. Sato K, Kawamura T, Wakusawa R (2000) Hepatic blood flow and function in elderly patients undergoing laparoscopic cholecystectomy. *Anesth Analg* 90: 1198–1202
261. Sauerland S, Neugebauer E (2000) Consensus conferences must include a systematic search and categorization of the evidence. *Surg Endosc* 14: 908–910
262. Saunders C, Battistella F, Whetzel T, Stokes R (1998) Percutaneous diagnostic peritoneal lavage using a Veress needle versus an open technique: A prospective randomized trial. *J Trauma* 44: 883–888
263. Schäfer M, Lauper M, Krähenbühl L (2001) Trocar and Veress needle injuries during laparoscopy. *Surg Endosc* 15: 275–280
264. Scheinin B, Kellokumpu I, Lindgren L, Haglund C, Rosenberg PH (1995) Effect of intraperitoneal bupivacaine on pain after laparoscopic cholecystectomy. *Acta Anaesthesiol Scand* 39: 195–198
265. Schilling MK, Redaelli C, Krähenbühl L, Signer C, Büchler MW (1997) Splanchnic microcirculatory changes during CO₂ laparoscopy. *J Am Coll Surg* 184: 378–383
266. Schindler E, Muller M, Kelm C (1995) Cerebral carbon dioxide embolism during laparoscopic cholecystectomy. *Anesth Analg* 81: 643–645
267. Schob OM, Alien DC, Benzel E, Curet MJ, Adams MS, Baldwin NG, Largiader F, Zucker KA (1996) A comparison of the

- pathophysiologic effects of carbon dioxide, nitrous oxide, and helium pneumoperitoneum on intracranial pressure. *Am J Surg* 172: 248–253
268. Schreiber JH (1990) Laparoscopic appendectomy in pregnancy. *Surg Endosc* 4: 100–102
 269. Schrenk P, Bettelheim P, Woisetschlager R, Rieger R, Wayand WU (1996) Metabolic responses after laparoscopic or open hernia repair. *Surg Endosc* 10: 628–632
 270. Schulte-Steinberg H, Weninger E, Jokisch D, Hofstetter B, Misera A, Lange V, Stein C (1995) Intraperitoneal versus interpleural morphine or bupivacaine for pain after laparoscopic cholecystectomy. *Anesthesiology* 82: 634–640
 271. Schulze S, Lyng KM, Bugge K, Perner A, Bendtsen A, Thorup J, Nielsen HJ, Rasmussen V, Rosenberg J (1999) Cardiovascular and respiratory changes and convalescence in laparoscopic colonic surgery: comparison between carbon dioxide pneumoperitoneum and gasless laparoscopy. *Arch Surg* 134: 1112–1118
 272. Schwenk W, Böhm B, Fugener A, Müller JM (1998) Intermittent pneumatic sequential compression (ISC) of the lower extremities prevents venous stasis during laparoscopic cholecystectomy. A prospective randomized study. *Surg Endosc* 12: 7–11
 273. Schwenk W, Böhm B, Junghans T, Hofmann H, Müller JM (1997) Intermittent sequential compression of the lower limbs prevents venous stasis in laparoscopic and conventional colorectal surgery. *Dis Colon Rectum* 40: 1056–1062
 274. Schwenk W, Böhm B, Witt C, Junghans T, Gründel K, Müller JM (1999) Pulmonary function following laparoscopic or conventional colorectal resection: a randomized controlled evaluation. *Arch Surg* 134: 6–13
 275. Schwenk W, Jacobi C, Mansmann U, Böhm B, Müller JM (2000) Inflammatory response after laparoscopic and conventional colorectal resections—results of a prospective randomized trial. *Langenbecks Arch Surg* 385: 2–9
 276. Schwenk W, Neudecker J, Mall J, Böhm B, Müller JM (2000) Prospective randomized blinded trial of pulmonary function, pain, and cosmetic results after laparoscopic vs microlaparoscopic cholecystectomy. *Surg Endosc* 14: 345–348
 277. Semm K, Arp WD, Trappe M, Kube D (1994) Schmerzreduzierung nach pelvi-/laparoskopischen Eingriffen durch Einblasen von körperwarmem CO₂-gas (Flow-Therme). *Geburtshilfe Frauenheilkunde* 54: 300–304
 278. Sezeur A, Bure-Rossier AM, Rio D, Savigny B, Tricot C, Martel P, Baubion O (1997) La coelioscopie augmente-t-elle le risque bactériologique de l'appendicéctomie? Résultats d'une étude prospective randomisée. *Ann Chir* 51: 243–247
 279. Shaneyfelt TM, Mayo-Smith MF, Rothwangl J (1999) Are guidelines following guidelines? The methodological quality of clinical practice guidelines in the peer-reviewed medical literature. *J Am Med Assoc* 281: 1900–1905
 280. Sharp JR, Pierson WP, Brady CE (1982) Comparison of CO₂- and N₂O-induced discomfort during peritoneoscopy under local anesthesia. *Gastroenterology* 82: 453–456
 281. Shea JA, Healey MJ, Berlin JA, Clarke JR, Malet PF, Starosick RN, Schwartz JS, Williams SV (1996) Mortality and complications associated with laparoscopic cholecystectomy. A meta analysis. *Ann Surg* 224: 609–620
 282. Shulman D, Aronson HB (1984) Capnography in the early diagnosis of carbon dioxide embolism during laparoscopy. *Can Anaesth Soc J* 31: 455–459
 283. Sietses C, Wierzer MJ, Eijbouts QAJ, Beelen RH, van Leeuwen PAM, von Blomberg BME, Meijer S, Cuesta MA (1999) A prospective randomized study of the systemic immune response after laparoscopic and conventional Nissen fundoplication. *Surgery* 126: 5–9
 284. Slim K, Bousquet J, Kwiatkowski F, Lescure G, Pezet D, Chipponi J (1999) Effect of CO₂ gas warming on pain after laparoscopic surgery: a randomized double-blind controlled trial. *Surg Endosc* 13: 1110–1114
 285. Smith RS, Fry WR, Morabito DJ, Koehler RH, Organ Jr CH (1995) Therapeutic laparoscopy in trauma. *Am J Surg* 170: 632–637
 286. Society of American Gastrointestinal Endoscopic Surgeons (1998) Guidelines for laparoscopic surgery during pregnancy. *Surg Endosc* 12: 189–190
 287. Soper NJ, Hunter JG, Petrie RH (1992) Laparoscopic cholecystectomy during pregnancy. *Surg Endosc* 6: 115–117
 288. Squirrell DM, Majeed AW, Troy G, Peacock JE, Nicholl JP, Johnson AG (1998) A randomized, prospective, blinded comparison of postoperative pain, metabolic response, and perceived health after laparoscopic and small incision cholecystectomy. *Surgery* 123: 485–495
 289. Stage JG, Schulze S, Moller P, Overgaard H, Andersen M, Rebsdorf-Pedersen VB, Nielsen HJ (1997) Prospective randomized study of laparoscopic versus open colonic resection for adenocarcinoma. *Br J Surg* 84: 391–396
 290. Stone J, Dyke L, Fritz P, Reigle M, Verrill H, Bhakta K, Boike G, Graham J, Gerbasi F (1998) Hemodynamic and hormonal changes during pneumoperitoneum and Trendelenburg positioning for operative gynecologic laparoscopy surgery. *Prim Care Update Ob Gyns* 5: 155
 291. Stuttmann R, Vogt C, Eypasch E, Doehn M (1995) Haemodynamic changes during laparoscopic cholecystectomy in the high-risk patient. *Endosc Surg Allied Technol* 3: 174–179
 292. Sungler P, Heinerman PM, Steiner H, Waclawiczek HW, Holzinger J, Mayer F, Heuberger A, Boeckl O (2000) Laparoscopic cholecystectomy and interventional endoscopy for gallstone complications during pregnancy. *Surg Endosc* 14: 267–271
 293. Szem JW, Hydo L, Barie PS (1996) A double-blinded evaluation of intraperitoneal bupivacaine vs saline for the reduction of postoperative pain and nausea after laparoscopic cholecystectomy. *Surg Endosc* 10: 44–48
 294. Tang C-L, Eu K-W, Tai B-C, Soh JGS, Machin D, Seow-Choen F (2001) Randomized clinical trial of the effect of open versus laparoscopically assisted colectomy on systemic immunity in patients with colorectal cancer. *Br J Surg* 88: 801–807
 295. Tang CS, Tsai LK, Lee TH, Su YC, Wu YJ, Chang CH, Tseng CK (1993) The hemodynamic and ventilatory effects between Trendelenburg and reverse Trendelenburg position during laparoscopy with CO₂-insufflation. *Ma Zui Xue Za Zhi* 31: 217–224
 296. Troidl H, Spangenberg W, Langen R, al-Jaziri A, Eypasch E, Neugebauer E, Dietrich J (1992) Laparoscopic cholecystectomy: technical performance, safety and patient's benefit. *Endoscopy* 24: 252–261
 297. Tsimoyiannis EC, Glantzounis G, Lekkas ET, Siakas P, Jabarin M, Tzourou H (1998) Intraperitoneal normal saline and bupivacaine infusion for reduction of postoperative pain after laparoscopic cholecystectomy. *Surg Laparosc Endosc* 8: 416–420
 298. Tsimoyiannis EC, Siakas P, Tassis A, Lekkas ET, Tzourou H, Kambili M (1998) Intraperitoneal normal saline infusion for postoperative pain after laparoscopic cholecystectomy. *World J Surg* 22: 824–828
 299. Unbehaun N, Feussner H, Slewert J (1995) Niederdruckinsufflationstechnik in der laparoskopischen Cholezystektomie. *Minimal Invasive Chirurgie* 4: 10–15
 300. Ure BM, Troidl H, Spangenberg W, Dietrich A, Lefering R, Neugebauer E (1994) Pain after laparoscopic cholecystectomy. Intensity and localization of pain and analysis of predictors in preoperative symptoms and intraoperative events. *Surg Endosc* 8: 90–96
 301. Ure BM, Troidl H, Spangenberg W, Neugebauer E, Lefering R, Ullmann K, Bende J (1993) Preincisional local anesthesia with bupivacaine and pain after laparoscopic cholecystectomy. A double-blind randomized clinical trial. *Surg Endosc* 7: 482–488
 302. Uzunkoy A, Coskun A, Akinci OF, Kocoyigit A (2000) Systemic stress responses after laparoscopic or open hernia repair. *Eur J Surg* 166: 467–471
 303. Vardulaki KA, Bennett-Loyd BD, Parfitt J, Normond C, Paisley S, Darzi A, Reeves BC (2000) A systematic review of the effectiveness and cost-effectiveness of laparoscopic surgery for colorectal cancer. A systematic review of the effectiveness and cost-effectiveness of laparoscopic surgery for colorectal cancer. *National Institute for Clinical Excellence, London*
 304. Vezakis A, Davides D, Gibson JS, Moore MR, Shah H, Larvin M, McMahon MJ (1999) Randomized comparison between low-pressure laparoscopic cholecystectomy and gasless laparoscopic cholecystectomy. *Surg Endosc* 13: 890–893
 305. Vik A, Jenssen BM, Brubakk AO (1991) Effect of aminophylline on transpulmonary passage of venous air emboli in pigs. *J Appl Physiol* 71: 1780–1786
 306. Volpino P, Cangemi V, D'Andrea N, Cangemi B, Piat G (1998) Hemodynamic and pulmonary changes during and after laparo-

- scopic cholecystectomy. A comparison with traditional surgery. *Surg Endosc* 12: 119–123
307. Walker CBJ, Bruce DM, Heys SD, Gough DB, Binnie NR, Eremin O (1999) Minimal modulation of lymphocyte and natural killer cell subsets following minimal access surgery. *Am J Surg* 177: 48–54
 308. Wallace DH, Serpell MG, Baxter JN, O'Dwyer PJ (1997) Randomized trial of different insufflation pressures for laparoscopic cholecystectomy. *Br J Surg* 84: 455–458
 309. Weber A, Muñoz J, Garteiz D, Cueto J (1997) Use of subdiaphragmatic bupivacaine instillation to control postoperative pain after laparoscopic surgery. *Surg Laparosc Endosc* 7: 6–8
 310. Wills VL, Hunt DR (2000) Pain after laparoscopic cholecystectomy. *Br J Surg* 87: 273–284
 311. Wills VL, Hunt DR, Armstrong A (2001) A randomized controlled trial assessing the effect of heated carbon dioxide for insufflation on pain and recovery after laparoscopic fundoplication. *Surg Endosc* 15: 166–170
 312. Wilson EB, Bass CS, Abrameit W, Roberson R, Smith RW (2001) Metoclopramide versus ondansetron in prophylaxis of nausea and vomiting for laparoscopic cholecystectomy. *Am J Surg* 181: 138–141
 313. Wittgen CM, Andrus CH, Fitzgerald SD, Baudendistel LJ, Dahms TE, Kaminski DL (1991) Analysis of the hemodynamic and ventilator effects of laparoscopic cholecystectomy. *Arch Surg* 126: 997–1001
 314. Wurst H, Schulte-Steinberg H, Finsterer U (1995) Zur Frage der CO₂-Speicherung bei laparoskopischer Cholezystektomie mit CO₂-Pneumoperitoneum. *Anaesthesist* 44: 147–153
 315. Yacoub OF, Cardona I, Coveler LA, Dodson MG (1982) Carbon dioxide embolism during laparoscopy. *Anesthesiology* 57: 533–535
 316. Yavuz Y, Ronning K, Lyng O, Marvik R, Gronbech JE (2001) Effect of increased intraabdominal pressure on cardiac output and tissue blood flow assessed by color-labeled microspheres in the pig. *Surg Endosc* 15: 149–155
 317. Yesildaglar N, Koninckx PR (2000) Adhesion formation in intubated rabbits increases with high insufflation pressure during endoscopic surgery. *Hum Reprod* 15: 687–691
 318. Yim SF, Yuen PM (2001) Randomized double-masked comparison of radially expanding access device and conventional cutting tip trocar in laparoscopy. *Obstet Gynecol* 97: 435–438
 319. Yoshida S, Ohta J, Yamasaki K, Kamei H, Harada Y, Yahara T, Kaibara A, Ozaki K, Tajiri T, Shirouzu K (2000) Effect of surgical stress on endogenous morphine and cytokine levels in the plasma after laparoscopic or open cholecystectomy. *Surg Endosc* 14: 137–140
 320. Yoshida T, Kobayashi E, Suminaga Y, Yamauchi H, Kai T, Toyama N, Kiyozaki H, Fujimura A, Miyata M (1997) Hormone-cytokine response. Pneumoperitoneum vs abdominal wall-lifting in laparoscopic cholecystectomy. *Surg Endosc* 11: 907–910
 321. Zieren J, Jacobi CA, Wenger FA, Volk HD, Müller JM (2000) Fundoplication: a model for immunologic aspects of laparoscopic and conventional surgery. *J Laparoendosc Adv Surg Tech A* 10: 35–40
 322. Zollinger A, Kraye S, Singer T, Seifert B, Heinzelmann M, Schlumpf R, Pasch T (1997) Haemodynamic effects of pneumoperitoneum in elderly patients with an increased cardiac risk. *Eur J Anaesthesiol* 14: 266–275
 323. Dietterle S, Pott C, Arnold A, Riedel HH (1998) The influence of insufflated carbon dioxide gas temperature on the pain intensity after pelviscopic procedures. *Min Invas Chir* 7: 103–105